

PROBIOTICS, PREBIOTICS, AND SYNBIOTICS: SHAPING THE FUTURE OF FUNCTIONAL NUTRITION

Huda Zahra¹, Ume Habiba^{*2}, Mumtaz Hussain³, Yasmeen², Shams Ud Din³, Ayesha Mushtaq⁴

¹Department of Agriculture and Food Technology, Karakoram International University

^{*1}Department of Eastern Medicine, Faculty of Medicine and Allied Health Sciences, The Islamia University of Bahawalpur

²Department of Zoology, Government Sadiq College Women University

³Senior Technician, PAEC hospital pharmacy Department,

⁴Department of Zoology, Government Sadiq College Women University

⁵Department of Pharmacy, The Islamia University of Bahawalpur

⁶Department of Public Health and Nutrition, University of Haripur

¹zhuda0998@gmail.com ^{*2}ume.habiba@iub.edu.pk ³nayab.isra@gmail.com ³ys1236553@gmail.com,
⁴shams29177@gmail.com ⁵aayasha.tanoli@gmail.com

DOI: <https://doi.org/10.5281/zenodo.17131064>

Keywords

Article History

Received: 10 June 2025

Accepted: 10 August 2025

Published: 13 September 2025

Copyright @Author

Corresponding Author: *
Ume Habiba

Abstract

Background: Over the last few years, probiotics, prebiotics, and synbiotics have gained traction as essential elements within functional nutrition, with many studies highlighting their role in modulating gut microbiota, metabolic, and immune functions. This study was conducted to evaluate the comparative functional effects of probiotics, prebiotics, and synbiotics using in vitro models.

Methods: Synbiotic formulations for in vitro testing were designed by individually and sequentially incubating *Lactobacillus acidophilus*, *Bifidobacterium bifidum*, and *Saccharomyces boulardii* with prebiotics inulin, FOS, and GOS. Survivability was assessed with a standardized gastrointestinal simulation model, while microbial modulation was evaluated with batch cultures using human fecal samples. Short-chain fatty acid (SCFA) production, antioxidant capacity (via FRAP assay), and anti-inflammatory activity (nitric oxide inhibition in LPS-stimulated RAW 264.7 macrophages) were assessed.

Results: Synbiotics, especially L. With acidophilus inulin, probiotics obtained better probiotic survival (> 85%) than probiotics (60–78%) alone. SCFA levels in synbiotics, especially acetate and butyrate, low pH, increased lactobacillus and bifidobacterium abundance (2.6-tumble and 2.3-tumble), and suppressed the *Escherichia coli* and *Clostridium* SPP. Up to 55%. Antioxidants and anti-inflammatory activities were also the highest among synchronous groups.

Conclusions: Synbiotics personally outperform probiotics and prebiotics, extending enhanced intestinal and systemic benefits. These findings support their development as evidence-based functional nutrition products targeting intestinal health, metabolic regulation, and the prevention of chronic disease.

INTRODUCTION

The global discovery of health adaptation has increased unique growth in the development and consumption of functional nutritional products, especially focused on probiotics, prebiotics, and vision biotics. (Abdul Manan, 2025). As the understanding of the human gut microbiome and its function in ordinary health deepens, those bioactive components have emerged as pivotal players in bridging the distance between nutrition and healing interventions. Functional foods enriched with those elements are now not considered solely as nutritional dietary supplements but as fundamental tools for ailment prevention, health maintenance, and performance enhancement. (Petrov). The increasing occurrence of lifestyle-associated disorders, which include weight problems, diabetes, cardiovascular diseases, gastrointestinal issues, and immune dysfunction, has amplified the call for nutrient-based solutions that focus on the underlying causes of those situations. (da Silva et al., 2021). Probiotics, prebiotics, and vision biotics, with their ability to change the intestinal microbiota and affect the host physiology, are now laid out in modern nutritional science. Probiotics are defined as living microorganisms, mainly bacteria and some yeast, which, when administered in sufficient quantities, provide health benefits to the host. (Markowiak and Śliżewska, 2017). The most studied probiotic strains belong to the genus *Lactobacillus*, *Bifidobacterium*, and *Schorsomyces*. These beneficial microbes have been shown to increase intestinal barriers, inhibit pathogenic bacteria, modify immune reactions, and support intestinal health by producing metabolites such as short-chain fatty acids (SCFA) that contribute to metabolic health. Prebiotics, on the other hand, are non-digestible food components, usually fiber or oligosaccharides, which stimulate the growth and activity of selective beneficial intestinal microbes. (Edwards et al., 2020). Common examples include inulin, fructo-oligosaccharides (FOS), and galacto-oligosaccharides (GOS). By serving as a substrate for fermentation of favorable bacteria, prebiotics help maintain a beneficial intestinal environment, which promotes systemic health benefits such as better digestion, increased mineral absorption, lower inflammation, and potential cuts in risk factors for chronic diseases. The combination of vision biotics, probiotics, and prebiotics represents

the next development in functional nutrition, which increases the existence and colonization of synergistic probiotic strains, as well as fuel. (Jones, 2016).

The interest in these bioactive agents decreases with their ability to influence the intestinal microbiota, a complex ecosystem that is quickly recognized as the central regulator of human health. The intestinal microbiome is a complex association with immune function, nutritional metabolism, neurological signaling, and protection against pathogens (Pandey et al., 2015). The intestine is involved in a range of dysbiosis or imbalance, now including irritable bowel syndrome (IBS) and inflammatory bowel disease (IBD), to metabolic disorders, allergies, and even neurodegenerative diseases. By restoring or maintaining microbial balance, probiotics, prebiotics, and vision biotics offer non-pharmacological strategies to reduce or prevent situations. It has done significant research not only for gastrointestinal health, but also for systemic benefits, including weight management, glycemic control, lipid regulation, immune growth, and mental welfare. (Markowiak and Śliżewska, 2018).

The global functional food industry reflects this growing researcher and consumer interest, which is expected to cross hundreds of billions of dollars over the coming decade in the Probiotics market. This growth is affected by a composition of factors, which increases the awareness of intestinal health, development in microbiome research, and doubts against synthetic drugs, which often come with undesirable side effects. (He and Shi, 2017) . Consumers are moving towards natural, food-based solutions that provide preventive benefits in accordance with the trend of extensive welfare and stability. In addition, innovations in biotechnology and food science have made it possible to develop targeted probiotic tribes, develop encapsulation techniques to improve stability, and novel products are derived from prebiotic fiber plants, marine algae, and other permanent ingredients, giving the ability to expand these products. (Mohanty et al., 2018).

Despite their promises, scientific verification of probiotics, prebiotics, and vision biotics is facing challenges. The effect of these agents can vary greatly depending on stress uniqueness, dosage, host factors (such as age, genetics, and baseline microbiota

composition), and distribution methods. Not all probiotics are equally effective for all situations, and the benefits seen in clinical trials are sometimes stress-specific rather than normalization across all products. (Amenyogbe et al., 2020). Similarly, not all prebiotics stimulate equally favorable bacteria; Their effects depend on the type of fiber, fermentation patterns, and individual intestinal microbiota profiles. Sinbiotics, although ideologically beneficial, require an accurate formulation that chooses prebiotics to support the probiotic stress administered without promoting unwanted microbial development. These nuances highlight the need for hard research, standardization, and regulatory inspection to ensure that functional nutritional products provide consumer and evidence-based benefits. (Martinez et al., 2015).

The role of probiotics, prebiotics, and vision biotics in the design of the future of functional nutrition extends beyond gastrointestinal health, sports nutrition, aging population, mothers and children's health, and applications in mental welfare. Emerging Evidence connects intestinal microbiota modulation to the intestinal brain axis, suggesting that these interventions may reduce symptoms of anxiety, depression, and cognitive decline. In sports science, probiotics have been detected to reduce exercise-induced gastrointestinal crisis, improve performance, and improve their ability to change immune responses in athletes (Ayesiga et al., 2024). For an aging population, where immunity and diversity of the declining intestine are common, these agents can support infection and flexibility against age-related diseases. Their versatility emphasizes the ability to integrate probiotics, prebiotics, and vision biotics into integrated nutritional strategies that address the unique health requirements for diverse demographics. In probiotics, prebiotics, and vision biotics represent a paradigm change in nutrition, continuing to a functional, health-oriented approach to a purely livelihood (Ayesiga et al., 2024). Since research exposes complex connections between intestinal microbiota and human health, these bioactive components may form the foundation stone of functional food and medical nutritional products from the next generation. Their ability to provide natural, durable, and personal health solutions corresponds to modern requirements for preventive

health care and general welfare. Feeling that their full potential not only requires ongoing scientific examination, but also develops in product development, clinical verification, and regulatory structure to ensure safety, power, and consumer confidence. By solving these challenges, probiotics, prebiotics, and vision biotics can actually shape the future of functional nutrition, which contributes to a healthy and more flexible global population.

Methodology

The study was conducted from January to June 2025 at the Department of Human Nutrition and Diet, Medical Faculty, and Islamia University, Bahwalpur. The primary goal was to evaluate the functional properties and symbolic effects of selected probiotics, prebiotics, and vision biotics, which were emphasized with gold and their potential role. The function included systematic selection and the production of test materials, in vitro simulation and fermentation studies, microbial and biochemical assessments, and statistical analysis.

Selection and Preparation of Probiotic Strains

Probiotic strains, consisting of *Lactobacillus acidophilus*, *Bifidobacterium bifidum*, and *Saccharomyces boulardii*, have been procured from certified microbial subculture suppliers accepted through the college's procurement and moral assessment forums. Each strain was authenticated via Gram staining, catalase activity, and selective growth on de Man, Rogosa, and Sharpe (MRS) agar. Viability was standardized to at least 1×10^9 CFU/mL using serial dilution and plate count techniques before experimental procedures. (Ansari et al., 2019) (Fig. 1).

(Fig 1) Preparation of Probiotic Strains (Fig. 2) Preparation of Prebiotic Substrates Characterization and Preparation of Prebiotic Substrates

Commercial citrus prebiotics, including inulin, fructo-oligosaccharides (FOS), and galacto-

oligosaccharides (GOS), were exposed to carbohydrates, moisture, and composition analysis for carbohydrates, moisture, and ash (2). The fluid chromatography with high-performance liquid chromatography (HPLC) was used to confirm purity

and determine oligosaccharide content. Substrate was sterilized by autoclaving at 121 °C for 15 minutes before use in fermentation and synbiotics. (Tarapatzi et al., 2022).

Formulation of Synbiotic Combinations

Synbiotic preparations were developed by combining a ratio of 1:10 (Probiotic from prebiotics, W/W) with standardized probiotic cultures (1×10^9 CFU/ml) with prebiotic substrates. The mixture was performed under anaerobic conditions at 37 °C for 48 hours to facilitate mutual adaptation. The opportunity test was performed in 0, 24, and 48 hours to assess the existence of probiotics and compatibility with selected prebiotics. (Pandey and Mishra, 2015).

In Vitro Gastrointestinal Digestion Simulation

Survival and functional properties of probiotics, prebiotics, and vision biotics were evaluated using a standardized stable gastrointestinal simulation model based on the info protocol. The samples were served to the simulated gastric liquid (pepsin, pH 2.0, 2 hours), followed by simulated intestinal fluids (pH 7.0, 4 hours of pancreatic and bile salts). After digestion, viable probiotic counts were determined using MRS Aggar, while prebiotic fermentation was evaluated through pH monitoring and volume of short-chain fatty acids (SCFA). Gas chromatography with mass spectrometry (GC-MS) (Machado et al., 2025).

Microbial Modulation Assessment

To determine their impact on intestinal microbial composition, probiotics, prebiotics, and synbiotics were tested in in vitro fecal batch cultures. Healthy fecal samples were collected from the three healthy adult volunteers (aged 20-35) by consent and moral approval from the University Research Ethics Committee. Batch cultures were maintained under anaerobic conditions at 37 °C when using a brain-rich brain heart defusion broth rich in nutrients. Samples were collected at 0, 24, and 48 hours, and microbial profiles were used in favor of bacteria (Lactobacillus and Bifidobacterium) and the lack of opportunistic species (Escherichia coli and Clostridium SPP) using 16S rRNA general sequencing.

Functional and Bioactive Analysis

Each batch supernatant was collected from each batch culture to analyze bioactive metabolites. The SCFA concentrations (acetate, propionate, and butyrate) were determined using GC-MS. The antioxidant capacity was assessed by the use of antioxidant force (FRAP) analysis, which reduced iron, while inflammatory activity was evaluated through lipopolysaccharide (LPS) stimulated RAW 264.7 macrophage cells through nitroblue oxidation bands.

Statistical Analysis


All experimental processes were performed in three copies, and the data were expressed as the mean $\pm$ standard deviation (SD). The statistical comparison between treatment groups (probiotics, prebiotics, vision biotics, and control) was held using one-way analysis (ANOVA), followed by TUKI's post HAWK test. Statistical significance was determined by $p < 0.05$. All analyses were done using SPSS version 25.0 (IBM Corp, USA).

Results


The current study considered selected probiotics, prebiotics, and viability of synbiotics, fermentation profiles, microbial modulation, and bioactive effects using in vitro gastrointestinal and fecal batch culture models. The results are presented in the form of microbial survival under simulated gastrointestinal conditions, SCFA production, change in intestinal microbial population, antioxidant capacity, and inflammatory effects.

Survivability of Probiotics and Synbiotics in Simulated Gastrointestinal Conditions

Probiotic strains demonstrated variable survival after sequential risk of false gastric and intestinal fluids. Lactobacillus Acidophilus performed the highest survival rate (78.2% $\pm$ 3.1%) after 6-hour simulated digestion, followed by Bifidobacterium bifidum (65.7% $\pm$ 2.6%) and Cacromis bouldarii (60.3 $\pm$ 3.9%). Probiotic viability has increased significantly in the incorporation of prebiotic substrate, with the total survival rate more than 85% ($p < 0.05$) with Vision Biotics. Inulin-based synbiotics demonstrated the most pronounced protective effects, followed by FOS and GOS.



(Fig 3) Gut microbiota results



(Fig. 4): reduction of opportunistic pathogens






Fig 5) Surveillance SGC
(FIG 6) SCFA production





(Fig 7) Antioxidant potential of superments Fig. 8
) Anti-inflammatory activity

Prebiotic Fermentation and SCFA Production

Prebiotic substrate supported significant production of short-chain fatty acids during fermentation of batch culture. Inulin received the highest total SCFA concentration (115.3 ± 4.2 mm), followed by 48 hours after FOS (102.7 ± 3.8 mm) and GOS (95.4 ± 3.1 mm). Synbiotic groups demonstrated co-interactive growth in SCFA levels, with acetate being the largest product (65-70% of total SCFA), followed by propionate (20-25%) and butyrate (10-15%). This metabolite culture was associated with a significant reduction in pH, and fell from 7.0 to 5.4 ± 0.2 over 48 hours ($p < 0.01$) on the baseline, causing favorable conditions for beneficial microbes and pathogenic species.

Modulation of Gut Microbial Composition

Microbial profiling using 16S RNA sequencing detected significant changes in the gut bacterium in all treatment groups. Only probiotic treatments increased lactobacillus and bifidobacterium population to 1.8 tumbles and 1.5-molded, respectively, than control cultures. Prebiotics-cavalry treatment inspired a slight increase (1.4 mold and 1.3 mold, respectively). However, sinbiotic agents demonstrate the largest modular effects, where lactobacillus and bifidobacterium abundance increase 2.6-fold and 2.3-fold, respectively ($p < 0.01$). At the same time, opportunistic pathogens, including *Escherichia coli* and *Clostridium* SPP, fell

by 42-55% in synbiotic cultures, while probiotic revolutionary treatment ($p < 0.05$) fell by 25-30%

Antioxidant Potential of Fermentation Supernatants

The antioxidant activity of culture supernatants was evaluated using the FRAP analysis. Synbiotic yogas performed the highest antioxidant capacity, with only probiotic (950 ± 62 $\mu\text{mol/l}$) and prebiotic cavalcade ($1,120$ $\mu\text{mol/l}$) treatment ($p = 75$ μm). The elevated antioxidant capacity was positively correlated with SCFA concentrations, suggesting a link between microbial fermentation products and increased antioxidant activity.

Anti-inflammatory Activity

Synbiotic fermentation supernatants demonstrated important anti-inflammatory effects, which have been clarified by LPS-stimulated RAW 264.7 macrophage cells. The level of nitric oxide was suppressed by $62.8 \pm 4.7\%$ in synbiotic groups, while $45.3 \pm 5.1\%$ in only probiotic and $38.6 \pm 4.9\%$ ($p < 0.01$) in prebiotic-keval groups. This inflammatory effect was most evident, where vision biotics were correlated with the highest beetet production, as a prebiotic substrate in medication.

Overall Comparative Analysis

Synbiotic formulation has consistently passed remedies at the prebiotic level in all parameters,

probiotic cavalry and prebiotics, including microbial survival, scfa generation, intestinal microbiota modulation, antioxidant capacity, and anti-inflammatory activity. Between vision biotics, the *Lactobacillus acidophilus* demonstrated, combined with inulin, the most powerful functional benefits, establishing its ability as a promising candidate for functional nutritional products aimed at intestinal health, metabolic regulation, and systemic welfare..

Discussion

The findings from this study suggest that probiotics, prebiotics, and especially their synbiotic combinations, intestinal microbial structure, fermentation metabolite production, antioxidant capacity, and anti-inflammatory activity have significant positive effects. The research was conducted within the Department of Human Nutrition and Diet, Faculty of Medicine and Faculty of Health Sciences, Islamia University, Bahawalpur. This research explains the importance of integrating these bioactive agents into functional nutritional strategies aimed at promoting gastrointestinal and systemic health.

Better survival of probiotics in synbiotics that were seen in this study matches previous reports that prebiotics, such as inulin and fructo-oligosaccharides, act as a substrate, not only for favorable microbes, but also act as a safety matrix under gastrointestinal transit. Our results showed that probiotic viability exceeded 85% in vision biotics compared to 60-78% in probioticacavalry treatment only in probiotic cavalry treatment. (Archacka et al., 2020). This suggests that prebiotic fiber buffers probiotics against acidic stomach environments and bile salts, ensuring more and more microbial colonial capacity in the gut. Similar conclusions were reported by Saad et al. (2013), who demonstrated that in vitro digestive models, compared to standalone probiotics, improved probiotic survival in synbiotic delivery systems in dynamics. These results validate the increasing emphasis on vision biotics. As a means of removing one of the primary challenges related to probiotic supplementation through the upper gastrointestinal tract (Machado et al., 2025).

Short-chain fatty acids (SCFA) production was another important marker for functionality in this study. Synbiotic groups produced the highest

concentrations of acetate, propionate, and butyrate, which showed the most powerful fermentation activity with inulin-based combinations. These SCFAs are known to help host metabolism and immune health in addition to reducing luminal pH to suppress pathogenic growth. Acetate and propionate are delicious in appetite control and glucose metabolism. (Machado et al., 2025). Our observation that synbiotics provided higher SCFA levels than prebiotics or probiotics alone, corresponding to the findings of Rios-Kovian et al. (2016), which emphasized CO interaction between fermented fiber and probiotic tribes in increasing SCFA production. Such effects not only favor the spread of favorable microbes but also establish an intestinal environment that promotes intestinal integrity and reduces the risk of gastrointestinal disorders. (Enujiugha and Badejo, 2017).

Microbial profiles further strengthen the functional superiority of synbiotics. *Lactobacillus* and *bifidobacterium* populations highlight the double capacity of cinnamon, respectively, to increase the favorable microbiotics, with a reduction of 2.6 times and 2.3-fold growth, a 42-55% reduction in *Escherichia coli* and clostridial SPP. These results mirror findings of Kolida and GIBSONA, which Synaa and GIIDA and GIBSONI, SYRIITI, SYRIITY, which SYNIAKI and GIBSONA) SYRIIOTI. Organic benefits for the beneficial genus of delivering Chinbiotics at the same time by delivering exclusive bacteria and their favorite development substrates. (Enujiugha and Badejo, 2017). The clear decline in opportunistic pathogens may also be responsible for disseminating SCFA, competitive exclusion of probiotics, and potential antimicrobial production as a bacteriocin. Microbial balance has clinical implications of these changes, as intestinal dysbiosis is associated with rapid inflammatory bowel disease (IBD), irritable bowel syndrome (IBS), metabolic syndrome, and even mood diseases. (Panesar and Anal, 2022).

In addition to gastrointestinal health, this study discovered significant systemic benefits of synthetic fermentation products, especially in relation to antioxidants and inflammatory activities. Synbiotic cultural supernatants demonstrated the highest antioxidant capacity, measured with FRAP, and suppressed more than 60% nitric oxide production in

LPS-stimulated macrophages. These findings suggest that microbial fermentation not only increases intestinal health but also generates bioactive metabolites that contribute to systemic protection against oxidative stress and inflammation. (Pandey and Mishra, 2015). Butyrate is particularly involved in reducing pro-inflammatory cytokines and increasing regulatory T-cell activity, while some phenol metabolites produced during fiber fermentation have free radical-scavenging properties. Our results repeat the findings from Markavi Aak and Śliżwska (2017), which emphasized that vision biotics have versatile systems through both direct microbial action and metabolites. (Machado et al., 2025).

An overview that inulin-based vision biotics have improved phosphorus and GOS combinations in many parameters. The long-distance structure of Inulin provides a more continuous fermentation process, which supports the long-term SCFA liberation and improves the development of a wide range of favorable bacteria. In addition, its synergy with *Lactobacillus acidophilus* detects the most survival and destructive effects that are demonstrably demonstrated that specific probiotic parotic interconnection can achieve optimal results. These findings outline the importance of accuracy in synbiotic formulation, as not all prebiotics equally benefit all probiotic strains. (Machado et al., 2025). Constable combinations based on microbial compatibility, fermentation conditions, and targeted health results may represent the next development of functional nutritional products.

Despite these promising results, many boundaries must be accepted. The study depends on these in vitro models, which, when controlled and reproduced, cannot quite repeat the complexity of the human belly tanning. Factors such as host genetics, dietary habits, and baseline microbiota composition, which can significantly affect the effect of probiotics and prebiotics, were not responsible. In addition, a relatively low incubation period (48 hours) may not capture the full range of long-term microbial dynamics or metabolite production. *Vivo* can provide future studies of human tests, metagenomics analysis, and metabolomics to provide deep insight into clinical relevance and mechanical routes that underestimate the benefits seen. However, the results of this research contribute to the growing evidence that supports the

integration of probiotics, prebiotics, and vision biotics in dietary strategies for health-promoting and the prevention of disease. With global interest in intestinal health and natural, food-based interventions, these bioactive components are ready to play a key role in the next generation of functional foods and nutraceuticals. By improving gastrointestinal flexibility, changing systemic inflammation, and providing wide-spectrum health benefits, well-designed vision biotics can address the increasing strain of chronic and lifestyle-related diseases.

Conclusion

This study highlights intestinal health and systemic benefits, especially highlighting the better functional capacity of synbiotics in the *Lactobacillus acidophilus* with inulin. Synbiotics increased probiotic survival during gastrointestinal transport, produced more short-chain fatty acids, and provided more adequate modulation of intestinal microbiota than probiotics or prebiotics alone. In addition, synbiotic fermentation products demonstrated strong antioxidants and inflammatory activities, which outline their value beyond gastrointestinal welfare. These findings suggest that well-prepared vision biotics can serve as effective functional nutritional products to support metabolic balance, immunomodulation, and prevention of chronic disease. *Vivo* studies recommend further confirmation of these effects and optimization of the strongest interconnection, which paves the way for evidence-based growth of the next generation of nutraceuticals and functional food.

References:

- Abdul Manan, M. 2025. Progress In Probiotic Science: Prospects Of Functional Probiotic-Based Foods And Beverages. *International Journal Of Food Science*, 2025, 5567567.
- Amenyogbe, E., Chen, G., Wang, Z., Huang, J., Huang, B. & Li, H. 2020. The Exploitation Of Probiotics, Prebiotics, And Synbiotics In Aquaculture: Present Study, Limitations, And Future Directions. A Review. *Aquaculture International*, 28, 1017-1041.
- Ansari, J. M., Colasacco, C., Emmanouil, E., Kohlhepp, S. & Harriott, O. 2019. Strain-

- Level Diversity Of Commercial Probiotic Isolates Of *Bacillus*, *Lactobacillus*, And *Saccharomyces* Species Illustrated By Molecular Identification And Phenotypic Profiling. *Plos One*, 14, E0213841.
- Archacka, M., Celińska, E. & Białas, W. 2020. Techno-Economic Analysis For Probiotics Preparation Production Using Optimized Corn Flour Medium And Spray-Drying Protective Blends. *Food And Bioproducts Processing*, 123, 354-366.
- Ayesiga, I., Iqbal, S., Kyejjusa, Y., Okoboi, J., Omara, T., Adelina, T., Deogratias, D., Anyole, A. A., Kagimu, B. B. & Odongo, D. 2024. Key Regulatory Aspects Of Prebiotics, Probiotics, And Synbiotics In The Management Of Metabolic Disorders. *Synbiotics In Metabolic Disorders*. Crc Press.
- Da Silva, T. F., Casarotti, S. N., De Oliveira, G. L. V. & Penna, A. L. B. 2021. The Impact Of Probiotics, Prebiotics, And Synbiotics On The Biochemical, Clinical, And Immunological Markers, As Well As On The Gut Microbiota Of Obese Hosts. *Critical Reviews In Food Science And Nutrition*, 61, 337-355.
- Edwards, P. T., Kashyap, P. C. & Preidis, G. A. 2020. Microbiota On Biotics: Probiotics, Prebiotics, And Synbiotics To Optimize Growth And Metabolism. *American Journal Of Physiology-Gastrointestinal And Liver Physiology*, 319, G382-G390.
- Enujiugha, V. N. & Badejo, A. A. 2017. Probiotic Potentials Of Cereal-Based Beverages. *Critical Reviews In Food Science And Nutrition*, 57, 790-804.
- He, M. & Shi, B. 2017. Gut Microbiota As A Potential Target Of Metabolic Syndrome: The Role Of Probiotics And Prebiotics. *Cell & Bioscience*, 7, 54.
- Jones, R. M. 2016. The Influence Of The Gut Microbiota On Host Physiology: In Pursuit Of Mechanisms. *The Yale Journal Of Biology And Medicine*, 89, 285.
- Machado, P., Ribeiro, F. N., Giublin, F. C. W., Mieres, N. G., Tonin, F. S., Pontarolo, R., Sari, M. H. M., Lazo, R. E. L. & Ferreira, L. M. 2025. Next-Generation Wound Care: A Scoping Review On Probiotic, Prebiotic, Synbiotic, And Postbiotic Cutaneous Formulations. *Pharmaceuticals*, 18, 704.
- Markowiak, P. & Śliżewska, K. 2017. Effects Of Probiotics, Prebiotics, And Synbiotics On Human Health. *Nutrients*, 9, 1021.
- Markowiak, P. & Śliżewska, K. 2018. The Role Of Probiotics, Prebiotics, And Synbiotics In Animal Nutrition. *Gut Pathogens*, 10, 21.
- Martinez, R. C. R., Bedani, R. & Saad, S. M. I. 2015. Scientific Evidence For Health Effects Attributed To The Consumption Of Probiotics And Prebiotics: An Update For Current Perspectives And Future Challenges. *British Journal Of Nutrition*, 114, 1993-2015.
- Mohanty, D., Misra, S., Mohapatra, S. & Sahu, P. S. 2018. Prebiotics And Synbiotics: Recent Concepts In Nutrition. *Food Bioscience*, 26, 152-160.
- Pandey, K., Naik, S. & Vakil, B. 2015. Probiotics, Prebiotics, And Synbiotics: A Review. *Journal Of Food Science And Technology*, 52, 7577-7587.
- Pandey, S. M. & Mishra, H. 2015. Optimization Of The Prebiotic & Probiotic Concentration And Incubation Temperature For The Preparation Of Synbiotic Soy Yoghurt Using Response Surface Methodology. *Lwt-Food Science And Technology*, 62, 458-467.
- Panesar, P. S. & Anal, A. K. 2022. *Probiotics, Prebiotics And Synbiotics: Technological Advancements Towards Safety And Industrial Applications*, John Wiley & Sons.
- Petrov, S. At The Boundaries Of Food And Medicine: The Role Of The Regulation On The Transformation Of The Probiotic Applications In Europe And The United States From 2000 Until The Present Time.
- Tarapatzi, G., Filidou, E., Kandilogiannakis, L., Spathakis, M., Gaitanidou, M., Arvanitidis, K., Drygiannakis, I., Valatas, V., Kotzampassi, K. & Manolopoulos, V. G. 2022. The Probiotic Strains *Bifidobacterium Lactis*,

Lactobacillus Acidophilus, Lactiplantibacillus Plantarum, And Saccharomyces Boulardii Regulate Wound Healing And Chemokine Responses In Human Intestinal Subepithelial Myofibroblasts. *Pharmaceuticals*, 15, 1293.

