



SYNTHESIS AND BIOLOGICAL EVALUATION OF TRANSITION METAL COMPLEXES OF PYRAZOLE SCHIFF BASE LIGAND

Sahrish Younus^{*1}, Tahir Maqbool², Amanullah Khan³, Areej Fatima⁴, Nida Younas⁵,
Sadia Younas⁴

^{*1}Department of Chemistry, University of Agriculture, Pakistan

²Department of Chemistry, Government College University Faisalabad, Pakistan

³Department of Veterinary Biomedical Sciences, Arid Agriculture University Rawalpindi, Pakistan

⁴Department of Allied Health Professionals, Government College University Faisalabad, Pakistan

⁵Department of Microbiology, Government College University Faisalabad, Pakistan

¹sahrish.m.younus@gmail.com

DOI: <https://doi.org/10.5281/zenodo.17200744>

Keywords

Pyrazole, Schiff base, antibacterial activity, Copper, azomethine, antifungal, Cadmium

Article History

Received on 04 July 2025

Accepted on 14 September 2025

Published on 25 September 2025

Copyright @Author

Corresponding Author: *

Sahrish Younus

Abstract


Pyrazole is the main component of bioactive agents, which act as antimicrobial, anti-inflammatory, analgesic, anticonvulsant, antibacterial, anxiolytic and antiviral agent. Pyrazole derivatives are also being used as therapeutic drugs which act on central nervous system (CNS). Azomethine derivatives of pyrazole also have biological potential and used as ligand to prepare metal complexes. Keeping the mentioned importance in view, in this research, we use amino pyrazole schiff base as ligand to prepare their metal complexes. The formation of products was monitored by using TLC, and the characterization of target compound has been done by melting point and FT-IR analysis. Antibacterial activity of these derivatives has also been studied.

INTRODUCTION


1.1. Pyrazole

Pyrazole is heterocyclic aromatic compound containing two adjacent nitrogen atoms in their ring with two double bonds and its structure is similar to imidazole (1), indazole (2), and


isoindazole (3) and some pyrazole derivatives also show non-aromaticity like pyrazolenine (4) isopyrazole (5) and 4,5-dihydro-1H-pyrazole (6) (Khan et al., 2016).



(1)



(2)



(3)



(4)



(5)



(6)

Figure 1.1: Some chemical structures of pyrazole derivatives



1.2. Drugs containing pyrazole ring

Pyrazole is heterocyclic compound which are used as medicinal drugs. For example, Indiplon (7) and Zaleplon (8) are drugs which contains pyrazole ring and act as anti-anxiety agent (Xu et al., 2012). CDPPB (9) is

used for the ministration of dementia and act as neuroleptic agent having pyrazole ring in their structure (Uslaner et al., 2009). Apixaban (10) is also a drug having a pyrazole ring and act as anticoagulant agent (Tripodi et al., 2015).




Figure 1.2: Drugs containing pyrazole ring

Diazepam (11) is important tranquilizer with a broad spectrum (Berghot et al., 2003). Similarly, pyrazole and pyrazoline derivatives act as analgesic and anti-erythrogenic agents (Bonesi et al., 2010). For example, Pentazocine (12) is a standard drug which contain pyrazole ring and act as analgesic agent (Vijesh

et al., 2013). Celecoxib (**13**) is a selective non-steroidal, anti-erythrogenic medicine which is used for the treatment of inflammation of osteoarthritis and acute pain in adults. Lonazolac (**14**) is also pyrazole containing non-steroidal anti-inflammatory drug. (Alka et al., 2011).




Figure 1.3: Derivatives of pyrazole (11), (12), (13) and (14) acts as pain-relieving and anti- inflammatory agents.

1.3. Pyrazole derivatives


Pyrazolo [3,4-d] pyrimidine (1,3-diazine) derivatives was act as antiviral, antimicrobial, meticidal, insecticidal, herbicidal and antileukemia agents (Abunada et al., 2008). Pyrazole derivatives are also being used as therapeutic drugs which act on the nervous system (CNS).

1.3.1. Antimicrobial activity


The fungal and bacterial infections are important threat and caused to death in immune deficiency persons (Vijesh et al., 2013). Many derivatives of multidrug are discovered which have ability against the microorganisms such as bacteria and fungi (Keche et al., 2012). For example, the compound (15) show the highest antimicrobial activity with better result against MRSA and QRSA strains (Song et al., 2013). The

synthesized compounds (16) having prazole ring in their molecule and most biologically active against *Streptococcus aureus*. (Shelke et al., 2012). Similarly, the newly prepared compound pyrazolopyrimidine (17) act as antibacterial agent with greater MIC against


S. aureus (Ghaneya et al., 2011). The compounds having NO₂ group at ortho positions in pyrazole ring are more active against *E. coli* bacteria (Alka et al., 2011).



(15)



(16)




(17)

Figure 1.4: Antimicrobial activities of pyrazole derivatives


1.3.2. Antimalarial activity

Malaria is a common disease which caused by *Plasmodium* genera. Female *Anopheles* mosquito spreads the malaria from one person to another. Mostly its target the underdeveloped countries like Pakistan and Africa due to high population density and poor infrastructure (Green et al., 2016). Many antimalarial drugs are prepared which having pyrazole ring and act as main nuclei against malarial disease. For examples, the compound (18) has five times


greater potency than chloroquine drug against malaria disease (Bekhit et al., 2015). Similarly, the prepared compound (19) is most active agent with higher antimalarial potency against chloroquine (Mowbray et al., 2009). The compound (20) is biologically active and act as schizonticidal and parasiticidal agent. Antimalarial activity of all these compounds is confirmed by *Plasmodium falciparum* (Alka et al., 2011).



(18)



(19)



(20)

Figure 1.5: Pyrazole derivatives containing antimalarial activity

1.3.3. Antihypertensive activity

Hypertension is characterized as a mulish medical condition in which blood in the arteries flow with more force than normal. Now-a-days, hypertension has

become a major disease throughout the world. There are many antihypertensive agents which contain pyrazole ring in their molecules (Lo et al., 2010). The newly prepared derivatives of pyrazole show

antihypertensive activity (21) which behave as more potent ACE inhibitor (Alka et al., 2011). The compound (22) containing fluorinated pyrazole which show the significant properties of antihypertensive

activity and compound (23) act as ACE inhibitors (Cui et al., 2013).




Figure 1.6: Pyrazole derivatives act as antihypertensive agents

1.3.4. Anticancer activity

Cancer is uncontrolled and serious life threatening disease in which abnormal cells are produced. Many scientists are trying to discover the best drugs for the treatment of cancer (Alka et al., 2011). Pyrazolopyrazinone derivatives mostly act as anticancer

agents (Newhouse et al., 2011). For example, the prepared compounds (24), (25) and (26) having pyrazole ring and show the anticancer activities against the apoptosis resistant cell lines HL-60 and K-562 (Jabali et al., 2011).




Figure 1.7: Anticancer activity of pyrazole derivatives

1.3.5. Antiviral activity

The infection caused by viruses is termed as viral infection. HCV virus damages the liver and cause the hepatitis C disease. Different pyrazole derivatives act as anti-viral agent (Xing et al., 2006). The newly prepared pyrazole containing compound (27) act as allosteric inhibitors in viruses (Koca et al., 2013). The

derivative of pyrazole (28) act as antiviral agent and prevent the eight diverse viral strains with an IC₅₀ of non-molar concentrations (Ashraf et al., 2018). Similarly, the pyrazole derivatives (29) show the antiviral activity against HIV (Wu et al., 2002).




Figure 1.8: Antiviral activity of pyrazole derivatives

1.3.6. Antifungal activity

Antifungal means kill fungi or stop the growth of fungi. Aspergilli is the main cause in safe traded off parasitic tainted patients (Vadiyala et al., 2017). Different five membered heterocyclic compounds such as pyrazole are very useful in medicinal field, mainly pyrazole have great value in the zone of antifungal agent (Foroumadi

et al., 2009). Similarly, compound (30) show antifungal activity (Samir et al., 2008) and (31) show antifungal activity against Fusarium Oxysporum (Ashraf et al., 2018) and the compound (32) exhibit in vitro antifungal activity against B. fabae and F. Oxysporum (Bondock et al., 2011).





Figure 1.9: Antifungal activity of pyrazole derivatives

1.3.7. Anti-aggravation activity

Aggravation is a multifactorial cycle that demonstrates the response of the living organism to different stimuli. Aggravation is characterized with numerous problems, for example, torment, redness of skin, growing, persistent aggravation may prompt joint inflammation, asthma, and psoriasis (Elena et al., 1999). Diverse pyrazole derivatives show anti-aggravation activity. For


example, Celecoxib (33) is a particular non-steroidal mitigating drug which is utilized for the treatment of acute pain and aggravation of osteoarthritis (Alka et al., 2011), (34) show anti- aggravation activity against carrageenan-incited rodent paw edema test (Aziz et al., 2009) and the compound (35) additionally show anti-aggravation activity (Ashish et al., 2014).

**Figure 1.10: Anti-inflammatory activity of pyrazole derivatives**

1.3.8. Antioxidant activity

Antioxidants are the compounds which interact and kill the free radical and prevent them from the cause of damage. Body additionally makes some free radical which is known as endogenous free radical which is influenced by age, diet and strength of a person. Free radicals are very reactive and they could damage our body because they affect the healthy cell of a molecule. Free radical can be dangerous for the body because

they damage the major component of the cell such as proteins, DNA and cell membrane (Rafat et al., 2014). Pyrazole is an important antioxidant because of the pyrazole ring which has important pharmacological and therapeutic properties. Such as pyrazole derivative (36), (37) which are analogue of coumarin derivatives (Aziz et al., 2009) and (38) also show antioxidant activity (Javarappa et al., 2012).

**Figure 1.11: Anti-oxidant activity of pyrazole derivatives**

1.3.9. Anti-Alzheimer's activity

Alzheimer disease is the cause of neurodegenerative disease which is 70% cause of dementia (Anh et al., 2019). Which is named as short-term loss of memory (Jabali et al., 2011)?

It normally happens gradually however as the infection increases then it became severe. Its normal signs are

issue with language, disposition swings, bewilderment and behavioral issues (Chris et al., 2014). Subsequently, new series of pyrazole derivative (39) which act as anti-alzheimer and the compound (40) also indicate property of alzheimer action (Valeria et al., 2013).




Figure 1.12: Anti-Alzheimer's activity of pyrazole derivatives

1.3.10. Anti-glaucoma activity

Glaucoma is a problem of eyes which effect the optic nerve. This illness may harm the eyes vision for all time if not treated it properly (Francesco et al., 2012). The main reason of glaucoma is visual hypertension for example the increased pressure within eyes. Glaucoma is predominantly treated by drug or either by medical procedure. There are different derivatives of pyrazole

which are used as anti-glaucoma agent (Rahmi et al., 2010). The pyrazole ring containing compound (41) and (42) show antiglaucoma activity. This showed that both of these compounds exhibit their inhibitory effects (Giselle et al., 2005). The pyrazole based compounds (43) and (44) also exhibit good anti-glaucoma activity (Rahmi et al., 2009).




Figure 1.13: Anti-glaucoma activity of pyrazole derivatives

1.4. Schiff base derivatives

Hugo schiff in 1864 reported azomethine derivatives which are commonly being known as schiff bases (Schiff et al., 1864). Azomethine derivatives can be synthesized by the reaction of essential amine with the aldehyde or ketones. Azomethine group is the

major part of these compounds whose general formula is $RCH=N-R_1$, here R and R₁ should be different substituents. Schiff base derivatives are usually biologically active compounds (Dhar et al., 1982; Przybyski et al., 2009).



For example; schiff base Pentazocine (45) is a standard drug containing of pyrazole ring act as analgesic agent and basic component of anthesia for surgery (Vijesh et al., 2013) and the compound (46) was studied against for multi-

drug-resistant bacteria (Ashraf et al., 2018). Amino pyrazole schiff base derivatives, for example, (47) show antitumor activity (Hassan et al., 2016).




Figure 14: Pyrazole Schiff base derivatives

1.4.1. Antimicrobial activity

Schiff bases such as (48), (49), and (50) show antimicrobial activity (Jabali et al., 2011).




Figure 15: Pyrazole Schiff base derivatives showing antimicrobial activity

1.4.2. Anti-bacterial activity

Schiff bases (51) and (52) synthesized from heterocyclic compound show antibacterial activity (Thakar et al., 2011), similarly

compound (53) was studied for their antibacterial activity against various strains of bacterium and parasite (Alka et al., 2011).





Figure 1.16: Antibacterial activity of pyrazole Schiff base derivatives


1.4.3. Antifungal activity

Different schiff bases obtained from different heterocyclic pyrazoles and triazoles show antifungal activity such as schiff base (54) is used as an antifungal agent (Sumangala et al., 2012),


schiff base (55) also show antifungal activity (Kouatli et al., 2010) and schiff base (56) show antifungal activity against different fungi (Kiran et al., 2012).



(54)



(55)



(56)

Figure 1.16: Antifungal activity of pyrazole schiff base derivatives

1.5. Pyrazole schiff base as a ligand

Pyrazole schiff bases are significant class of ligand in coordination chemistry and have significant application in the field of science such as antifungal, antimalarial, antineoplastic and antiviral agent (Monica et al., 2001). Schiff bases are also used as chelating agents in coordination chemistry when they have

stabilizing groups such as -OH close to -HC=N group. The ligand schiff base are also important in analytical chemistry for determination of metal (Praveen et al., 2007). For example, compound (57) is used as a ligand to from palladium complexes and exhibit



1.6. Metal complexes of pyrazole schiff base and their biological importance

Pyrazole schiff bases from metal complex upon coordination with different metals. Schiff base derivative of pyrazole has been employed for complex formation. Schiff base derivatives are commonly used as anti-pyretic, anti-microbial, anti-fungal, anti-inflammatory, anti-malarial, anti-fungal, antiproliferative and antibacterial agents. A coordination complex having metal atom in center is called a metal complex (Nevin et al 2011). Metal complexes of pyrazole along

1.6.1. Antitumor and Anticancer activity

The compound (63) shows cytotoxic activity which is more than of Cisplatin which is an

with its derivatives usually used as herbicides, fungicides and in drugs. For example, new schiff base (60) from complexes that show antibacterial activity against *Staphylococcus* bacteria (kiran et al., 2012) and pyrazole-based ligand (61) also from complexes (Susmita et al., 2016). Similarly, compound (62) these are used as powerful antifungal, antibacterial and antiviral agents (Giselle et al., 2005).

anticancer drug (Abu et al., 2006) and the compound (64) show antitumor and anticancer activity (Ribeiro et al., 2008).




Figure 1.20: Metal complexes of Pyrazole Schiff base

1.6.2. Anti-bacterial activity

To study the biological significance of schiff base, the pyrazole schiff bases and their metal complex (65) have been examined for their antibacterial action (Kiran et al., 2012). The metal complex (66) possess insulin-mimetic activity and they can be used for storage and

detoxification purpose (Aziz et al., 2009) and the metal complex (67) has tried for their antibacterial action against *Escherichia coli* and made to interact with DNA to judge on their binding capacity by absorption (Anupama et al., 2012).




Figure 1.22: Metal complexes (65) show antibacterial, (66) show insulin-mimetic and (67) also have antibacterial activity

1.7. Furan derivatives

The presence of furan ring is much important in natural products which exhibit the biologically activity (Hou et al., 1998; Lipshutz, 1986). Many pharmaceutical products are obtained from substituted aryl furan-2-carbaldehyde (Dunlop et al., 1953). Furfural (2-furaldehyde) is heteroaromatic compound having aldehyde functional group. Commercially, 2-

furaldehyde is synthesized from dehydration of pentosans (polysaccharides) in the presence of acid catalysis by using continuous reactor. In the commercial production of 2-furfuraldehyde, large amount of agricultural wastes is used for this purpose which including almond husks, oat hulls, corncobs etc. (Dias et al., 2010).

2-furfuraldehyde is mostly used for cleansing of diesel fuels as well as lubricating fuels and many solvents

such as 2-furancarbinol, 2-methyloxolane, adipic ketone (cyclopentanone) and THF (Yan et al., 2014), 4-valerolactone (GV L) (Bui et al., 2013). Furfural (2-furfuraldehyde) is also used for the preparation of 2-carboxyfuran, 2-Furanmethanol (Zeynizadeh et al., 2005) or 2-furylcyanide (Furan-2-carbonitrile) intermediates (Ghaneya et al., 2011). Furan resins are prepared from furfuryl alcohol, the most important furan derivatives such as 2-carboxyfuran, 2-methyl

furan, 4-oxopentanoic acid and furfuryl amine (Zeitsch, 2000).

The furan derivatives are used as medicines like Dantralene (68) Clodanolene (69) and Azimilide (70) due to high biological activity. Dentrolene (Dentamacrine) and Clodanolene are standard drugs which containing furan ring and used for the treatment of muscle (muscle relaxant) (Mashkovskii et al., 2000). Azimilide is drug which having furan ring and show the activity against heart diseases (Bui et al., 2013).




Figure 1.23: Drugs containing furan moiety

The furan derivatives such as 2,5-disubstituted furan are synthesized via cross coupling of palladium based catalyzed reactions of metalated furans. The preparation of 5-aryl-furan-2-carbaldehyde carried out using cross coupling reactions in which palladium used as catalyst with zero oxidation state (Balachari et al., 2000).

The papulacandin derivatives are prepared by substituted Furfuran derivatives and Papulacandins D are act as antifungal agent. The Papulacandin families are biological active and act as anti-microbial agents. They help in the treatment of are *P. carinii* pneumonia and control the AIDS (Schmatz et al., 1990).


Mentioning the importance of furan, pyrazole and metal complexes, we have planned to

introduce furan derivatives on to 1-phenyl-3-methyl-5-aminopyrazole in order to from schiff bases along with their metal complexes. The aims and objectives of the studies are

- To prepare suitable substituted aryl furfural.
- To prepare schiff base derivatives and their metal complexes using substituted pyrazole and furfural derivatives.
- To characterize all the target compounds through different techniques like M.P, TLC and FTIR.
- To evaluate the target compounds for their antibacterial activity.

3.1. Methods

3.1.1. Synthesis of 5-(4-nitrophenyl)-2-furaldehyde (71)



4-Nitroaniline (0.2g, 5 mmol) was dissolved in the mixture of H₂O and concentrated HCl, stirred the mixture 15 minutes followed by heating until the formation of a clear solution. The reaction mixture was cooled in ice bath up to -5 °C. After cooling the reaction mixture, a chilled solution of NaNO₂ (0.25g, 5.1 mmol) in H₂O (5 mL) was added portion wise into the reaction mixture with constant stirring and kept the temperature below 0 °C. The reaction mixture after the addition of sodium nitrite was left for one hour at 0 °C for completion of diazotization. Then to the solution of 2-furaldehyde (0.2g, 2.5mmol) in five ml water and five ml acetone diazonium solution was poured drop wise followed by cupric chloride in water. The temperature of the solution was raised up to 30 °C and the reaction mixture was stirred for 4-6 hours and left for 24 hours. The product obtained on filter paper was dried and recrystallized from suitable solvent (Aslam et al., 2013).

Yield: 55%

M.P: 190-192 °C (Literature, 196 °C) (Aslam et al., 2013)


Solubility: Ethyl acetate, methyl alcohol, ethyl alcohol

TLC: n-hexane and Ethyl acetate solution (1:1)

FT-IR: 2750 cm⁻¹ (C-H of formyl), 1666.65 cm⁻¹ (C=O), 1513.75 and 1326.30 cm⁻¹(NO₂),

3.2. Synthesis of Pyrazole Schiff base ligands (72-73)

3.2.1. General Procedure






R= H, 4-NO₂-Ph

Pyrazole schiff base were synthesized through reaction of 1-phenyl-3-methyl-5- amino pyrazole (0.2g, 0.00086 mol) with furfural derivatives (100 mg, 1mmol) in AcOH using 6ml. The reaction mixture was refluxed for five hours. Then this reaction mixture was cooled for ten minutes at room temperature and 6ml of distilled water was added into that reaction mixture to form precipitates. These precipitates were washed with distilled H₂O and ETOH and it was recrystallized using ethyl alcohol.

3.2.2. Synthesis of N-(furan-2-yl-methylene)-4-methyl-1-phenyl-1H-pyrazole-3- amine (L1) (72)



Yield: 67%


M.P: 180 °C

Solubility: Ethyl acetate, methyl alcohol, chloroform and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1597 cm⁻¹ (C=N), 3117.59 cm⁻¹ (aromatic stretching), 2923.70 cm⁻¹ (C-H stretching), 1503.9 cm⁻¹ shows (C=C), and 1259 cm⁻¹ (C-O) and 1335 cm⁻¹ (C-N).

3.2.3. Synthesis of 4methyl-N-(5-(4-nitrophenyl) furan-2-yl-methylene)-1- phenyl-1H-pyrazole-3- amine (L2) (73)



Yield: 77%

M.P: 190 °C


Solubility: Ethyl acetate, methyl alcohol, chloroform and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1601 cm⁻¹(C=N), 3377.38 cm⁻¹ (aromatic stretching), 2915.56 cm⁻¹ (C-H stretching), 1537.84 cm⁻¹ (C=C), 1508.67 cm⁻¹ & 1333.84 cm⁻¹(NO₂), 1203 cm⁻¹ (C-O) and 1333 cm⁻¹ (C-N).


3.3. Synthesis of metal complexes of Pyrazole (74-80)

3.3.1. General Procedure



Pyrazole schiff base were synthesized through reaction of 1-phenyl-3-methyl-5-amino pyrazole (0.2g, 0.00086 mol) with furfural derivatives (100 mg, 1mmol) in AcOH using 6ml. The reaction mixture was refluxed for five hours. Then this reaction mixture was cooled for ten minutes at room temperature and 6ml of distilled water was added into that reaction mixture to form precipitates. These precipitates were washed with distilled H₂O and ETOH and it was recrystallized using ethyl alcohol.

3.3.2. Synthesis of N-(furan-2-yl-methylene)-4-methyl-1-phenyl-1H-pyrazo-3- amine (L1) (72)



Yield: 77%

M.P: 190 °C

Solubility: Ethyl acetate, methyl alcohol, chloroform and ethyl alcohol


TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1601 cm⁻¹(C=N), 3377.38 cm⁻¹ (aromatic stretching), 2915.56 cm⁻¹ (C-H stretching), 1537.84 cm⁻¹ (C=C), 1508.67 cm⁻¹ & 1333.84 cm⁻¹(NO₂), 1203 cm⁻¹ (C-O) and 1333 cm⁻¹ (C-N).



3.4. Synthesis of metal complexes of Pyrazole (74-80)

General Procedure



Metal complexes were synthesized by using pyrazole schiff base as ligands. The ligand (0.2g, 0.00086 mol) was dissolved in 5ml of ethanol in a beaker and metal chloride (0.1g, 0.00043) dissolved in 7ml of ethanol in a round bottom flask. The ligand solution was poured into the metal solution and reflux the reaction 4 hours. Precipitates obtained were filtered off and washed several times with distilled H₂O and EtOH. It was finally recrystallized by using ethanol.

3.4.1. Synthesis of Cu²⁺ complex using L1 (74)

Yield: 60%

M.P: 305 °C

Solubility: Ethyl acetate, methyl alcohol and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1541.66 cm⁻¹(C=N), 2360.66 cm⁻¹ (aromatic stretching), 1507.68 cm⁻¹(C=C), 1319.69 cm⁻¹(C-N) and 490.67 cm⁻¹ (M-N).

3.4.2. Synthesis of Cu²⁺ complex using L2 (75)

Yield: 67%

M.P: 300 °C

Solubility: Ethyl acetate, methyl alcohol and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1596 cm⁻¹(C=N), 2360.66 cm⁻¹ (aromatic stretching), 3032.45 cm⁻¹ (OH stretching), 1457.06 cm⁻¹ (C=C), and 488.96 cm⁻¹ (M-N).

3.4.3. Synthesis of Ni²⁺ complex using ligand L2 (76)

Yield: 60%

M.P: 275 °C

Solubility: Ethyl acetate, methyl alcohol and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1599 cm^{-1} (C=N), 3336.11 cm^{-1} (OH), 2361 cm^{-1} (aromatic stretching), 1456.30 cm^{-1} (C=C) and 491 cm^{-1} (M-N).

3.4.4. Synthesis of Cd²⁺ complex using ligand L2 (77)

Yield: 69%

M.P:300 $^{\circ}\text{C}$

Solubility: Ethyl acetate, methyl alcohol and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1599.17 cm^{-1} (C=N), 2368.19 cm^{-1} (aromatic stretching), 1457.14 cm^{-1} (C=C), 491.42 cm^{-1} (M-N) and 544.22 cm^{-1} (M-O).

3.4.5. Synthesis of Zn²⁺ complex using ligand L2 (78)

Yield: 70%

M.P: 270 $^{\circ}\text{C}$

Solubility: Ethyl acetate, methyl alcohol and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1598.48 cm^{-1} (C=N), 2360.31 cm^{-1} (aromatic stretching), 1456.97 cm^{-1} (C=C), 1275.86 cm^{-1} (C-N) and 492 cm^{-1} (M-N).

3.4.6. Synthesis of Fe²⁺ complex using L2 (79)

Yield: 60%

M.P: \geq 250 $^{\circ}\text{C}$

Solubility: Ethyl acetate, methyl alcohol and ethyl alcohol

TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1598 cm^{-1} (C=N), 3115.71 cm^{-1} (OH), 2361.39 cm^{-1} (aromatic stretching), 1474.60 cm^{-1} (C=C), 1275 cm^{-1} (C-N) 500 cm^{-1} (M-N).

3.4.7. Synthesis of Co²⁺ complex using ligand L2 (80)

Yield: 73%

M.P: 290 $^{\circ}\text{C}$

Solubility: Ethyl acetate, methyl alcohol and ethyl alcohol



TLC: n-hexane and Ethyl acetate Solution (3:1)

FT-IR: 1599 cm⁻¹ (C=N), 3245.75 cm⁻¹ (OH), 2363.55 cm⁻¹ (aromatic stretching), 1454.62 cm⁻¹ and 495 cm⁻¹ (M-N).


RESULTS AND DISCUSSION

Arylation of furfural to prepare its new derivative is important due to medicinal uses of furan derivatives. For examples, Nitrafudan, Clodanolene and Azimilide have aryl furan moiety in their structures (Obushak et al., 2009). 4-Nitrofurfural furan derivative was synthesized through copper catalyzed Meerwein arylation in which arenediazonium salt is intermediate. Similarly, pyrazole are one of naturally dynamic compounds utilized in drugs (Elena et al., 1999). These are antiviral, antibacterial, antifungal and antitumor (Shah et al., 2011 & Satish et al., 2008). So, keeping in mind the importance of both compounds some new derivative of pyrazole schiff bases were synthesized by the reaction of pyrazole with furfural derivatives and by utilizing these heteroaryl schiff base different type of metal complexes were synthesized. The formation of derivatives was monitored through thin layer chromatography. The melting point of given compounds was also checked. The structural composition of synthesized compounds was done through FT-IR.

4.1. Synthesis of substituted aryl furfural derivative

4.1.1. Synthesis of 5-(4-Nitrophenyl) furan-2-carbaldehyde (71)

NO₂
H
(71)



Scheme 4.1: Synthesis of 5-(4-Nitrophenyl) furan-2-carbaldehyde (71)

5-(4-Nitrophenyl) furan-2-carbaldehyde was synthesized through the Meerwein arylation by using 4-Nitro anilines and furfurals. The completion of reaction and purity of its product was checked by TLC and melting point. The value of melting point of newly arylated product was also correlated with the value of literature. The FT-IR spectra of arylated furfural showed the peak for carbonyl group at 1666.68 cm⁻¹ and nitro (Nitro group) showed sharp two peaks at 1513.32 and 1326.79 cm⁻¹. The FT-IR of the compound (71) is given next.





Figure 4.1: FT-IR spectrum of 5-(4-nitrophenyl) furan-2-carbaldehyde



Mechanism of arylation of furfural




Scheme 4.2: Mechanism of arylation of furfural

4.2. Synthesis of pyrazole schiff base ligands (72-73)

4.2.1. Synthesis of N-(furan-2-yl-methylene)-4-methyl-1-phenyl-1H-pyrazo-3-amine L1 (72)

4.2.2.



Scheme 4.4: Synthesis of N-(furan-2-yl-methylene)-4-methyl-1-phenyl-1H-pyrazo-3-amine L1 (72)

Pyrazole schiff base was prepared by the reaction of 1-phenyl-3-methyl-5-amino pyrazole with furfural in acetic acid 6ml. The reaction mixture was refluxed for four hours. The purity of the compound was confirmed through thin layer chromatography and further characterization was done through FT-IR. The melting point of compound (72) was 180°C. No signal was observed for carbonyl (C=O) and amino group in the FT-IR spectra of the product which indicate that it is converted into desired product. In spectra peak at 1597cm⁻¹ is the peak of (C=N) which show synthesis of schiff base, peak at 3117.59 cm⁻¹ shows (aromatic stretching), peak at 2923.70 cm⁻¹ shows (C-H stretching), and peak at 1503.9 cm⁻¹ shows (C=C), and 1259 cm⁻¹ show (C=O) and 1335 cm⁻¹ show (C-N) group (Alpana et al., 2009). The FT-IR of (72) is given below.





Figure 4.2: FT-IR spectrum of N-(furan-2-yl-methylene)-4-methyl-1-phenyl-1H-pyrazo-3-amine L1 (72)

Synthesis of 4-methyl-N-(5-(4-nitrophenyl) furan-2-yl-methylene)-1- phenyl-1H-pyrazole-3-amine L2 (73)


Scheme 4.5: Synthesis of 4-methyl-N-(5-(4-nitrophenyl) furan-2-yl-methylene)-1-phenyl-1H-pyrazole-3-amine L2 (73)
 Pyrazole schiff base has prepared by the reaction of 1-phenyl-3-methyl-5-amino pyrazole with 5-(4-nitrophenyl)-2-furaldehyde in acetic acid 6ml. The reaction mixture was refluxed for five hours. Thin layer chromatography has confirmed purity of compound and furthermore characterization was done through the FT-IR. The melting point of compound (73) was 190°C and the percentage yield of the reaction was 67%. In FT-IR spectra the peak at 1601 cm⁻¹ is of (C=N) which indicate the formation of schiff base. No signal was observed for carbonyl (C=O) and amino group in the FT-IR of spectra of the compound which indicate that it is converted into desired product. Along with other peaks at 3377.38 cm⁻¹ show (aromatic stretching), peak at 2915.56 cm⁻¹ show (C-H stretching), and show 1537.84 cm⁻¹ show (C=C), (NO₂) shows two peaks at 1333.84 cm⁻¹ & 1508.67 cm⁻¹, peak at 1203 cm⁻¹ show (C-O) and 1333 cm⁻¹ indicate the presence of (C-N) group (Arun et al., 2008) . The FT-IR spectrum of compound (73) is given below.




Figure 4.3: FT-IR spectrum of 4-methyl-N-(5-(4-nitrophenyl) furan-2-yl-methylene)-1-phenyl-1H-pyrazole-3-amine L2 (73)

4.3. Synthesis of metal complexes pyrazole schiff base (74-80)

4.3.1. General procedure

Metal complexes were synthesized by using pyrazole Schiff base as ligands. The ligand (0.2g, 0.86 mmol) was dissolved in 5ml of ethanol in a beaker and metal chloride (0.43mmol) dissolved in 7ml of ethanol in a round bottom flask. The ligand solution was poured into the metal solution and reflux the reaction 4 hours. Precipitates formed were filtered off and washed several times with distilled H₂O and EtOH. It was finally recrystallized by using ethanol.



Scheme 4.7: Synthesis of Cu²⁺ complex of L1 (74)

Metal [Cu(II)] complex of ligand L1 is synthesized with the reaction of ligand with metal salt of copper (CuCl₂.2H₂O) in 2:1 using EtOH as solvent. The melting point of the compound

(74) was 305 °C. The percentage yield of compound was 60%. The purity of the compound was checked with TLC. In the FT-IR spectra showed the peak at 1541.66 cm⁻¹ which is assigned to (C=N). The peak of (C=N) in case of ligand is observed at 1597 cm⁻¹. The shifting towards lower frequency indicate that the ligand is involved in coordination with metal, peak at 1507.68 cm⁻¹ show (C=C) , peak at 1319.69 cm⁻¹ show (C-N) bond, 2360.66 cm⁻¹ show (aromatic stretching) and peak at 490.67 cm⁻¹ show (M-N) group which show metal coordination bond (Shehab et al., 2004). The FT-IR spectrum of (74) is given below.





Figure 4.4: FT-IR spectrum of Cu²⁺ metal complex of L1 (74)

4.3.2. Synthesis of Cu²⁺ complex of ligand L2 (75)



Scheme 4.8: Synthesis of Cu²⁺ complex of L2 (75)



Metal [Cu(II)] complex of ligand L2 is prepared with the reaction of ligand with metal salt of copper ($\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$) in 2:1 using EtOH. The melting point of compound (75) was 300°C . The percentage yield of compound was 67%. The purity of the compound was checked by TLC. In the FT-IR spectra showed the peak at 1596 cm^{-1} which is assigned to (C=N). The peak of

(C=N) in case of ligand is observed at 1601 cm^{-1} . The shifting towards lower frequency indicate that the ligand is involved in coordination with metal, peak at 2360.66 cm^{-1} show 'aromatic stretching', peak at 1457.06 cm^{-1} show (C=C), and peaks at 488.96 cm^{-1} shows the existence of (M-N) bond (Kiran et al., 2012). The FT-IR spectrum of (75) given below.





Figure 4.5: FT-IR spectrum of Cu^{2+} metal complex of L2 (75)

4.3.3. Synthesis of Ni^{2+} complex of L2 (76)



Scheme 4.9: Synthesis of Ni^{2+} complex of L2 (76)

Metal [Ni(II)] complex of ligand L2 is prepared by the reaction of ligand with metal salt of nickel ($\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$) in 2:1 using EtOH. The melting point of compound (76) was 275°C . The percentage yield of compound was 60%. The purity of the compound was checked with TLC. In the FT-IR spectra showed the peak at 1599 cm^{-1} which is assigned to (C=N). The peak of (C=N) in case of ligand is observed at 1601 cm^{-1} . The shifting towards lower frequency indicate that the ligand is involved in coordination with metal, and further peak at 3336.11 cm^{-1} show the presence of (OH) group, 2361 cm^{-1} (aromatic stretching), peak at 1456.30 cm^{-1} show (C=C), and 491 cm^{-1} indicate (M-N) metal coordination bond (Abd et al., 2011). The FT-IR spectrum of (76) is given below.





Figure 4.5: FT-IR spectrum of Ni^{2+} metal complex of L2 (76)

4.3.4. Synthesis of Cd²⁺ complex of L2 (77)





Scheme 4.10: Synthesis of Cd²⁺ complex of L2 (77)

Metal [Cd(II)] complex of ligand L2 is synthesized with the reaction of ligand with metal salt of cadmium (CdCl₂.H₂O) in 2:1 using EtOH. The melting point of compound (77) was 300°C. The percentage yield of compound was 69%. The purity of the compound was checked by TLC. In the FT-IR spectra showed the peak at 1599.17 cm⁻¹ which is given to (C=N). The peak of (C=N) in case of ligand is observed at 1601 cm⁻¹. The shifting towards lower frequency indicate that the ligand is involved in coordination with metal, peak at 2361 cm⁻¹ show (aromatic stretching) and further peak at 1457.14 cm⁻¹ show (C=C), peak at 491.42 cm⁻¹ show (M-N) group and peak at 544.22 cm⁻¹ show (M-O) group (Saugata et al., 2011). The FT-IR spectrum of (77) is given below.





Figure 4.6: FT-IR spectrum of Cd²⁺ metal complex of L2 (77)

4.3.5. Synthesis of Zn²⁺ complex of L2 (78)





Scheme 4.11: Synthesis of Zn^{2+} complex of L2 (78)

Metal $[Zn(II)]$ complex of ligand L2 is formed by the reaction of ligand with metal salt of Zinc ($ZnCl_2$) in 2:1 using EtOH. The melting point of compound (78) was $275^{\circ}C$. The percentage yield of compound was 70%. The purity of the compound was checked with TLC. In the FT-IR spectra showed the peak at 1598.48 cm^{-1} which is given to (C=N). The peak of (C=N) in case of ligand is observed at 1601 cm^{-1} . The shifting towards lower frequency indicate that the ligand is involved in coordination with metal, peak at 2360.31 cm^{-1} show (aromatic stretching), peak at 1456.97 cm^{-1} show (C=C), peak at 1275.86 cm^{-1} show (C-N) and 492 cm^{-1} show the presence of (M-N) group (Kiran et al., 2012). The FT-IR spectrum of (78) is given below.





Figure 4.8: FT-IR spectrum of Zn^{2+} metal complex of L2 (78)

4.3.6. Synthesis of Fe^{2+} complex of L2 (79)





Scheme 4.12: Synthesis of Fe^{2+} complex of L2 (79)

Metal [Fe(II)] complex of ligand L2 is prepared by the reaction of ligand with metal salt of iron (FeCl₂) in 2:1 using EtOH as solvent. The melting point of compound (79) was ≥ 250 °C. The percentage yield of compound was 60%. The purity of the compound was checked with TLC. In the FT-IR spectra showed the peak at 1598 cm⁻¹ which is given to (C=N). The peak of (C=N) in case of ligand is observed at 1601 cm⁻¹. The shifting towards lower frequency indicate that the ligand is involved in coordination with metal, peak at 3115.71 cm⁻¹ show (OH) group, 2361.39 cm⁻¹ show (aromatic stretching) peak at 1474.60 cm⁻¹ show (C=C), peak at 1275 cm⁻¹ show (C-N) bond and further peak at 500 cm⁻¹ indicate (M-N) group (Saugata et al., 2011). The FT-IR spectrum of (79) is given below.





Figure 4.9: FT-IR spectrum of Fe^{2+} metal complex of L2 (79)

4.3.7. Synthesis of Co^{2+} complex of L2 (80)





Scheme 4.13: Synthesis of Co^{2+} complex of L2 (80)

Metal [Co(II)] complex of ligand L2 is prepared with the reaction of ligand with metal salt of cobalt (CoCl_2) in EtOH. The melting point of compound (80) was 290°C . The percentage yield was compound is 73%. The purity of the compound was examined by TLC. In the FT-IR spectra showed the peak at 1599 cm^{-1} which is given to ($\text{C}=\text{N}$). The peak of ($\text{C}=\text{N}$) in case of ligand is observed at 1601 cm^{-1} . The shifting towards lower frequency indicate that the ligand is involved in coordination with metal, 3245.75 cm^{-1} show (OH) group, peak at 1454.62 cm^{-1} show ($\text{C}=\text{C}$), 2363.55 cm^{-1} show (aromatic stretching) and 495 cm^{-1} show (M-N) bond (Kiran et al., 2012). The FT-IR spectrum of (80) is given below.




Figure 4.11: FT-IR spectrum of Co^{2+} metal complex of L2 (80)

Physical properties of arylated furfural derivatives are given in table 4.1.

Table 4.1: Physical properties of arylated furfural derivative, pyrazole schiff bases and their metal complexes (71-80)

Compound	Molecular formula	Mol. Mass (g/mol)	Colour and physical state	Solubility	Yield (%)	M.P ($^{\circ}\text{C}$)
71	$\text{C}_{11}\text{H}_7\text{NO}_4$	217	Yellow (powder)	EtOH, MeOH, EtOAc	55	190-192
72	$\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}$	251	Black (powder)	EtOH, MeOH, EtOAc	67	180
73	$\text{C}_{21}\text{H}_{16}\text{N}_4\text{O}_3$	372	Pale yellow (powder)	EtOH, MeOH, EtOAc	77	190
74		566	Light brown	EtOH, MeOH,	60	305



	²⁺ C ₃₀ H ₂₆ CuN ₆ O ₂		(powder)	EtOAc		
75	²⁺ C ₄₂ H ₃₂ CuN ₈ O ₆	808	Dark red (powder)	EtOH, MeOH, EtOAc	67	300
76	²⁺ C ₄₂ H ₃₂ NiN ₈ O ₆ ²⁺	803	Yellow (powder)	EtOH, MeOH, EtOAc	60	275
77	²⁺ C ₄₂ H ₃₂ CdN ₈ O ₆	857	Light green (powder)	EtOH, MeOH, EtOAc	69	300
78	²⁺ C ₄₂ H ₃₂ NO ₆ Zn ²⁺	808	Dark yellow (powder)	EtOH, MeOH, EtOAc	70	270
79	²⁺ C ₄₂ H ₃₂ FeN ₈ O ₆ ²⁺	801	Brown (powder)	EtOH, MeOH, EtOAc	60	≥250
80	²⁺ C ₄₂ H ₃₂ CoN ₈ O ₆	804	Dark yellow (powder)	EtOH, MeOH, EtOAc	73	290

Table 4.2: FT-IR spectral values for arylated furfural derivatives, pyrazole schiff bases and their metal complexes (71-80)

Compound number	FT-IR
71	2750 cm ⁻¹ (C-H of formyl), 1666.65 cm ⁻¹ (C=O), 1513.75 and 1326.30 cm ⁻¹ (NO ₂).
72	1597 cm ⁻¹ (C=N), 3117.59 cm ⁻¹ (aromatic stretching), 2357.87 cm ⁻¹ (aromatic ring), 2923.70 cm ⁻¹ (C-H stretching), 1503.91 & 1455.60 cm ⁻¹ show (C=C), and 1259 cm ⁻¹ (C-O) and 1335 cm ⁻¹ (C-N).
73	1601 cm ⁻¹ (C=N), 3377.38 cm ⁻¹ (aromatic stretching), 2915.56 cm ⁻¹ (C-H stretching) 2357.87 cm ⁻¹ (Aromatic ring), 1537.84 cm ⁻¹ (C=C), 1508.67 cm ⁻¹ & 1333.84 cm ⁻¹ (NO ₂), 1203 cm ⁻¹ (C-O) and 1333 cm ⁻¹ (C-N).
74	1541.66 cm ⁻¹ (C=N) and 490.67 cm ⁻¹ (M-N).



75	1596 cm ⁻¹ (C=N), and 488.96 cm ⁻¹ (M-N).
76	1599 cm ⁻¹ (C=N), 3336.11 cm ⁻¹ (OH) and 491 cm ⁻¹ (M-N).
77	1599.17 cm ⁻¹ (C=N), 491.42 cm ⁻¹ (M-N) and 544.22 cm ⁻¹ (M-O).
78	1598.48 cm ⁻¹ (C=N) and 492 cm ⁻¹ (M-N).
79	1598 cm ⁻¹ (C=N), 3115.71 cm ⁻¹ (OH) and 500 cm ⁻¹ (M-N).
80	1599 cm ⁻¹ (C=N), 3245.75 cm ⁻¹ (OH) and 495 cm ⁻¹ (M-N).

Anti-bacterial activity

The prepared derivatives of furan based pyrazole Schiff base derivatives along with their metal complexes were dissolved in suitable solvent and subjected to anti-bacterial activity by disk diffusion method. These derivatives were tested against bacterial strain by using gentamycin as positive control. The zone of inhibition of all the synthesized compounds were measured in millimeter (mm) against *E. coli* (anti-bacterial) strains.

All these derivatives showed zone of inhibition 8.5-11.5 mm as compare to reference gentamycin 21mm (for antibacterial). The results of some selected pyrazole schiff base derivatives and their metal complexes showed the moderate anti-bacterial activity because these derivatives showed lower values of zone of inhibition against two strains as compare to reference gentamycin as shown in figures and table.




Figure : Zone of inhibitions of compounds against *E.coli* (Front)



Figure : Zone of inhibitions of compounds against *E. coli* (Back)

Table: Antimicrobial activity of synthesized compounds


Compound number	Concentration 40mg/ml	Zone of inhibition (mm) Bacteria <i>S. nigrum</i>
Gentamycin	+ev	17mm
73	10mg/ml	12mm
77	20mg/ml	14.5mm
78	30mg/ml	16mm
75	40mg/ml	18.5mm

SUMMARY

Different arylated furfurals compounds were prepared by Meerwein arylation. Pyrazole schiff bases were prepared from furfural derivatives. By using pyrazole schiff different kind of metal complexes were prepared. The purity of synthesized products has been determined by TLC and characterization of prepared compounds (products) was done via melting point and FT-IR. The spectra of prepared derivatives show sharp peaks for the

presence of different substituents. Some selected derivatives were assessed for antimicrobial activities, among the following series compound (75) showed the most highly antimicrobial activity.

Following arylated furfural derivative, pyrazole schiff bases and their metal complexes were prepared (71-80), which are given below.



References

Abbas Rahmati, Miranda Eskandari-Vashareh, Meysam Alizadeh-Kouzehrash (2013). Synthesis of 3-(benzylidene amino)-2-phenyl-5H-imidazo[1,2-b] pyrazole-7-carbonitriles via a four-component condensation reaction. *Tetrahedron*, 69:4199-4204.

Abd, El-Motaleb M. Ramadan (2011). Macroyclic nickel(II) complexes: synthesis, characterization, superoxide scavenging activity and DNA-binding. *Journal of Molecular Structure*, 1015:56-66.

Abunada, N., Hassaneen, H., Kandile, N. & Miqdad, O. (2008). Synthesis and antimicrobial activity of some new pyrazole, fused pyrazolo [3, 4-d] pyrimidine and pyrazolo [4, 3-e] [1, 2, 4] triazolo [1, 5-c] pyrimidine derivatives. *Molecules*, 13(7): 1501-1517.

Abu-Surrah, Adnan S. Kettunen, Mike (2006). Platinum group antitumor chemistry design and development of new anticancer drugs complementary to Cisplatin. *Current Medicinal Chemistry*, 13:1337-1357.

Adnan A. Bekhit, Manal N. Saudi, Ahmed M.M. Hassan, Salwa M. Fahmy, Tamer M. Ibrahim, Doaa Ghareeb, Aya M. El-Seidy, Sherry N. Nasralla, Alaa El-Din A. Bekhit (2009). Synthesis, in silico experiments and biological evaluation of 1,3,4-trisubstituted pyrazole derivatives as antimalarial agents. *European Journal of Medicinal Chemistry*, 163: 353-366.

Adnan S. Abu-Surrah, Kayed A. Abu Safieh, Iman M. Ahmad, Maher Y. Abdalla, Mikdad T. Ayoub, Abdussalam K. Qaroush, Ahmad M. Abu-Mahtheieh (2019). New palladium(II) complexes bearing pyrazole-based Schiff base ligands: Synthesis, characterization and cytotoxicity. *European Journal of Medicinal Chemistry*, 45:471-475.

Alka Chauhan, P. K. Sharma, Niranjan Kaushik (2011). Pyrazole: A Versatile Moiety. *International Journal of ChemTech Research*, 3(1): 11-17.

Alpana K Gupta and Rajendra S Sindal. (2009). A comparative study of electrochemical reduction of isatin its synthesized Schiff bases at HMD. *Journal of Chemical Sciences*, 121: 347-351.

Andressa Esteves Souza, Aurea Echevarro, Ivo Vencato, Maria Luisa Jimenoc and Jose Elguero (2001). Unexpected formation of bis-pyrazolyl derivatives by solid support coupled with microwave irradiation. *Tetrahedron*, 57: 6147-6153.

Anh P. Truong, Danielle L. Aubele, Gary D. Probst, Martin L. Neitzel, Chris M. Semko, Simeon Bowers et al (2019). Design, synthesis, and structure-activity relationship of novel orally efficacious pyrazole/sulfonamide based dihydroquinoline Y-secretase inhibitors. *Bioorganic & Medicinal Chemistry*, 19:4920-4923.

Anita Sharma and Manish Shah (2013). Synthesis and Characterization of some transition metal complexes derived from bidentate schiff base ligand. *Journal of Applied Chemistry*, 3(5): 62-66.

Anupama Kamath, Krishna Naik, Sandeep P. Netalkar, Dhoolesh, Gangaram Kokare, Vidyanand

K. Revankar (2012). Pyrazole-bridged late first row transition metal complexes derived from hexadentate compartmental ligand: synthesis, characterization, antibacterial activity, and DNA binding/cleavage studies. *Medicinal Chemistry Research*, 22: 1948-1956.

Arun M. Isloor, Balakrishna Kalluraya, Prashanth Shetty (2008). Regioselective reaction: Synthesis, characterization and pharmacological studies of some new Mannich bases derived from 1,2,4-triazoles. *European Journal of Medicinal Chemistry*, 44: 3784-3787.

Ashish Kumar Tewari, Ved Prakash Singh, Pratima Yadav, Garima Gupta, Amit Singh, Raj Kumar Goel, Pravin Shinde, C. Gopi Mohan (2014). Synthesis, biological evaluation and molecular modeling study of pyrazole derivatives as selective COX-2 inhibitors and anti-inflammatory agents. *Bioorganic Chemistry*, 56:8-15.

Ashraf S. Hassan, Gaber O. Moustafa, Ahmed A. Askar, Ahmed M. Naglah and Mohamed A. Al- Omar (2018). Synthesis and antibacterial evaluation of fused pyrazole and Schiff bases, *Synthetic Communication*, 48(21): 2761-2772.

Ashraf S. Hassan, Hanem M. Awad. Asmaa A. Maged-El-Din. Taghrid S. Hafez. (2017). Synthesis and in vitro antitumor evaluation of novel Schiff bases. *Medecinal Chemistry Research*, 27: 915-927.

Aslam, S., Asif, N., Khan, M. N., Khan, M. A., Munawar, M. A. & Nasrullah, M. (2013). Synthesis of novel arylfurfuryl chalcones. *Asian Journal of Chemistry*, 25(14): 7738-7742

Aziz MA, Rahma Gedaa, Hassan AA. (2009). Synthesis of novel pyrazole derivatives and evaluation of their antidepressant and anticonvulsant activities. *European Journal of Medicinal Chemistry*, 44: 3480-3487.

Balachari, D. & O'Doherty, G. A. (2000). Sharpless asymmetric dihydroxylation of 5-aryl-2-vinylfurans: application to the synthesis of the spiroketal moiety of papulacandin D. *Organic Letters*, 2(6): 863-866.

Bekhit, A. A., Hassan, A. M., El Razik, H. A. A., El-Miligy, M. M., El-Agroudy, E. J. & Bekhit, A. E. D. A. (2015). New heterocyclic hybrids of pyrazole and its bioisosteres: design, synthesis and biological evaluation as dual acting antimalarial-antileishmanial agents. *European Journal of Medicinal Chemistry*, 94: 30-44.

Berghot, M. A. & Moawad, E. B. (2003). Convergent synthesis and antibacterial activity of pyrazole and pyrazoline derivatives of diazepam. *European Journal of Pharmaceutical Sciences*, 20(2): 173-179.

Bondock S, Khalifa W, Fadda AA (2011). Synthesis and antimicrobial activity of some new 4-hetarylpyrazole and furo [2, 3-c] pyrazole derivatives. *European Journal of Medicinal Chemistry*, 46:2555-2561.

Bonesi, M., Loizzo, M. R., Statti, G. A., Michel, S., Tillequin, F. & Menichini, F. (2010). The synthesis and angiotensin converting enzyme (ACE) inhibitory activity of chalcones and their pyrazole derivatives. *Bioorganic & Medicinal Chemistry Letters*, 20(6): 1990-1993.

Bui, L., Luo, H., Gunther, W. R. & Roman-Leshkov, Y. (2013). Domino reaction catalyzed by zeolites with Brønsted and Lewis acid sites for the production of γ -valerolactone from furfural. *Angewandte Chemie International Edition*, 52(31): 8022-8025.

Chris G. Whiteley (2014). Arginine metabolizing enzymes as targets against Alzheimers' disease. *Neurochemistry International*, 67:23-31.

Cui, J. J. (2013). Targeting receptor tyrosine kinase MET in cancer: small molecule inhibitors and clinical progress. *Journal of Medicinal Chemistry*, 57(11): 4427-4453.

Dhar, D.N. & Taploo, C.L. (1982). Schiff-bases and their applications. *Journal of Scientific & Industrial Research*, 41(8): 501-506.

Dias, T., Bronze, M. R., Houghton, P. J., Mota-Filipe, H. & Paulo, A. (2010). The flavonoid-rich fraction of Coreopsis tinctoria promotes glucose tolerance regain through pancreatic function recovery in streptozotocin-induced glucose-intolerant rats. *Journal of Ethnopharmacology*, 132(2): 483-490.

Dunlop, A. P. & Peters, F. N. (1953). The Furans (Vol. 1). Reinhold Pub. Corp, Elinson, M. N., Dorofeev, A. S., Nasybullin, R. F. & Nikishin, G. I. (2008). Facile and convenient synthesis of 4, 4'-(arylmethylene) bis (1H-pyrazol-5-ols) by electrocatalytic tandem Knoevenagel- Michael reaction. *Synthesis*, 2008(12): 1933-1937.

Elena I, Klimovo, Marcos Martinez Garcia, Tatiana Klimovo Berestneva, Cecilio Alvarez Toledano, Ruben Alfredo Toscano, Lena Ruiz Ramirez (1999). The structure of bicyclic ferrocenyl methylene substituted 2-pyrazolines and their reactions with azodicarboxylic acid N-phenylimide. *J. Organomet. Chemistry*, 585: 106-111.

Foroumadi A, Sorkhi M, Moshafi, Safavi M, Rineh AR, Shafiee A, Emami S (2009). 2-Substituted-5-nitroheterocycles: In vitro antihelicobacter pylori activity and structure-activity relationship study. *Medicinal Chemistry*, 5:529-534.

Francesco. Di, M. E., Avolio, S., Pompei, M., Pesci, S., Monteagudo, E., Pucci, V. & Summa, V. (2012). Synthesis and antiviral properties of novel 7-heterocyclic substituted 7-deazaadenine nucleoside inhibitors of Hepatitis C NS5B polymerase. *Bioorganic & Medicinal Chemistry*, 20(15): 4801-4811.

Ghaneya Sayed Hassan, Hanan Hassan Kadry, Sahar Mahmoud Abou-Seri, Mamdouh Moawad Ali c, Abeer Essam El-Din Mahmoud (2011). Synthesis and in vitro cytotoxic activity of novel pyrazolo[3,4-d] pyrimidines and related pyrazole hydrazones toward breast adenocarcinoma MCF-7 cell line. *Bioorganic & Medicinal Chemistry*, 19: 6808-6817.

Giselle Cerchiaro, Katia Aquilano, Giuseppe Filomeni, Giuseppe Rotilo, Maria Rosa Ciriolo, Ana Maria Da Costa Ferreria. (2005). Isatin schiff base copper complexes and their influence on cellular viability. *Journal of Inorganic Biochemistry*, 99: 1433-1440.

Green, J. L., Moon, R. W., Whalley, D., Bowyer, P. W., Wallace, C., Rochani, A. & Ansell, K. H. (2016). Imidazopyridazine inhibitors of *Plasmodium falciparum* calcium-dependent protein kinase 1 also target cyclic GMP-dependent protein kinase and heat shock protein 90 to kill the parasite at different stages of intracellular development. *Antimicrobial Agents and Chemotherapy*, 60(3): 1464-1475.

Hassan AS, Hafez TS, Ali M. M, Khatab TK. (2016). Design, synthesis and cytotoxic activity of some new pyrazolines bearing benzofuran and pyrazole moieties. *Research Journal Pharmaceutical Biological Chemistry Science*, 7: 417-429.

Hou, X. L., Cheung, H. Y., Hon, T. Y., Kwan, P. L., Lo, T. H., Tong, S. Y. & Wong, H. N. (1998).

Regioselective syntheses of substituted furans. *Tetrahedron*, 54(10): 1955-2020. J.C. Pessoa, I. Correia (2017). New Cu(II) complexes with pyrazolyl derived Schiff base ligands synthesis and biological evaluation. *Journal of Inorganic Biochemistry*. 174:63- 75.

Jabali J. Vora, Akash R. Patel, Dinesh R. Patel, Sandip Dholakiya (2011). Synthesis and antimicrobial activity of some new (E)-4-chloro-2-((3-substituted aryl)-1-phenyl-1H-pyrazole-4-yl)methylene amino)-6-methoxy pyrimidine-5-carbaldehyde derivatives. *American Journal of Pharmtech Research*, 3(1): 2249-3387.

Javarappa Rangaswamy, Honnaiah Vijay Kumar, Salakatte Thammaiah Harini, Nagaraja Naik (2012). Synthesis of benzofuran based 1,3,5-substituted pyrazole derivatives: As a new class of potent antioxidants and antimicrobials- A novel accost to amend biocompatibility. *Bioorganic & Medicinal Chemistry Letters*, 22:4773-4777.

Keche, A. P., Hatnapure, G. D., Tale, R. H., Rode, A. H. & Kamble, V. M. (2012). Synthesis, anti-inflammatory and antimicrobial evaluation of novel 1-acetyl-3, 5-diaryl-4, 5-dihydro (1H) pyrazole derivatives bearing urea, thiourea and sulfonamide moieties. *Bioorganic & Medicinal Chemistry Letters*, 22(21): 6611-6615.

Khan, M. F., Alam, M. M., Verma, G., Akhtar, W., Akhter, M. & Shaquiquzzaman, M. (2016). The therapeutic voyage of pyrazole and its analogs: a review. *European Journal of Medicinal Chemistry*, 120: 170-201

Kiran Singh, Yogender Kumar, Parvesh Puri, Mahender Kumar, Chetan Sharma. (2012). Cobalt, nickel, Copper and Zinc Complexes with 1,3-diphenyl-1H-pyrazole-4-carboxaldehyde schiff bases: Antimicrobial, Spectroscopic, Thermal and fluorescence studies. *European Journal of Medicinal Chemistry*, 52: 313-321.

Koca, I., Özgür, A., Coşkun, K. A. & Tutar, Y. (2013). Synthesis and anticancer activity of acyl thioureas bearing pyrazole moiety. *Bioorganic & Medicinal Chemistry*, 21(13): 3859- 3865.

Kouatli Omar, Athina Geronikaki, Panagiotis Zoumpoulakis, Charalabos Camoutsis, Marina Sokovic, Ana Ceric, Jasmina Glamocilja (2010). Novel 4-thiazolidinone derivatives as potential antifungal and antibacterial drugs. *Bioorganic & Medicinal Chemistry*, 18:426-432.

Lipshutz, B. H. (1986). Five-membered heteroaromatic rings as intermediates in organic synthesis. *Chemical Reviews*, 86(5): 795-819.

Lo, H. Y., Man, C. C., Fleck, R. W., Farrow, N. A., Ingraham, R. H., Kukulka, A. & Sharma, R. (2010). Substituted pyrazoles as novel SEH antagonist: investigation of key binding interactions within the catalytic domain. *Bioorganic & Medicinal Chemistry Letters*, 20(22): 6379-6383.

Manohare S.V, and S. S. Thakare (2019). Synthesis and antibacterial screening of Schiff base derived from 3-(5-bromo thiopen-yl)-1-(4-chloro phen-2-yl)-1H-pyrazole-4- carbaldehyde. Manohare and Thakare, 10(8): 3741-3745.

Mashkovskii, M. D. (2000). *Lekarstvennye sredstva (drugs)*. Moscow, Novaya Volna, 2, 295.

Mohamed Rifaat Hamza Elmoghayer, Mohamed Ali Elsayed Khalifa, Mohamed Kamal Ahmed Ibraheim, and Mohamed Hilmy Elnagdi. (1981). Reactions with heterocyclic β -enaminoesters; A novel synthesis of 2-amino-3-ethoxycarbonyl-(4H)-pyrans. *Monatshefte fur Chemie*, 113: 53-57.

Monica M.B. Pessoa, Gustavo F.S. Andrade, Vera R. Paoli Monteiro, Marcia L.A. Temperini (2001). 2-Formyl pyridine thiosemicarbazone and methyl derivatives spectroscopic studies. *Polyhedron*, 20:3133-3141.

Mowbray, C. E., Burt, C., Corbau, R., Perros, M., Tran, I., Stupple, P. A. & Wood, A. (2009). Pyrazole NNRTIs 1: Design and initial optimization of a novel template. *Bioorganic & Medicinal Chemistry Letters*, 19(19): 5599-5602.

Nadia Ribeiro, Roberto E. Di Paolo, Adelino M. Galvao, Fernanda Marques (2018). Photophysical properties and biological evaluation of a Zinc(II)-5-methyl-1H-pyrazole Schiff base complex. *Spectrochemical Acta Part A: Molecular and Biomolecular Spectroscopy*, 204:317-327.

Nevin Turan, Esin Kaya, Bayram Gunduz, Naki Colak, and Hanifi Korkoca. (2011). Synthesis, characterization of poly (E)-3-amino-((3-bromo phenyl) diaziny)-1H-pyrazol-5-ol: Investigation of antibacterial activity, Fluorescence, and optical properties. *Fibers and Polymers*, 13: 415-424.

Newhouse, B. J., Hansen, J. D., Grina, J., Welch, M., Topalov, G., Littman, N. & Brandhuber, B. J. (2011). Non-oxime pyrazole based inhibitors of B-Raf kinase. *Bioorganic & Medicinal Chemistry Letters*, 21(11): 3488-3492.

Praveen. A Kumar, P. Raveendra Reddy, and V. Krishna Reddy (2007). Direct and derivative spectrophotometric determination of Cobalt (II) in microgram quantities with 2-hydroxy- 3-methoxy benzaldehyde thiosemicarbazone. *Journal of the Korean Chemical Society*, 51:331.

Przybylski, P., Huczynski, A., Pyta, K., Brzezinski, B. & Bartl, F. (2009). Biological properties of Schiff bases and azo derivatives of phenols. *Current Organic Chemistry*, 13(2): 124-148.

Rafat M. Mohareb, Amira E. M. Abdallah, Mahmoud A. Abdelaziz (2014). New approaches for the synthesis of pyrazole, thiophene, thieno [2,3-b] pyridine, and thiazole derivatives together with their anti-tumor evaluations. *Medicinal Chemistry Research*, 23: 564-579.

Rahmi Kasımcıoğlu, Metin Bulbul, B. Seçkin Arslan, Basak Gökçe (2010). Synthesis, characterization and antiglaucoma activity of some novel pyrazole derivatives of 5-amino-1,3,4-thiadiazole-2-sulfonamide. *European Journal of Medicinal Chemistry*, 45:4769-4773.

Rahmi Kasimogullari, Metin Bulbul, Hatice Gunhan, Hulya Guleryuz (2009). Effects of new 5-amino-1,3,4-thiadiazole-2-sulfonamide derivatives on human carbonic anhydrase isozymes. *Bioorganic & Medicinal Chemistry*, 17:3295-3301.

Ribeiro, N. S. Roy, N. Butenko, I. Cavaco, T. Pinheiro, I. Alho, F. Marques, F. Avecilla, N. Satish Rai, Balakrishna Kalluraya, B. Lingapa, Shalini Shaenoy, Vagdevi, G. Puranic (2008). Convenient access to 1,3,4-trisubstituted pyrazoles carrying 5-nitrothiophene moiety via 1,3-dipolar cycloaddition of sydnone with acetylenic ketones and their antimicrobial evaluation. *European Journal of Medicinal Chemistry*, 43: 1715-1720.

Rodolfo Moreno-Fuquen, Fernando Cuenu, John Eduard Torres, Gala De la Vega, Esperanza Galarza, Rodrigo Abonia, Alan R. Kennedy (2017). Presence of $\pi\ldots\pi$ and $\text{C-H}\ldots\pi$ interactions in the new Schiff base 2-[(E)-[(3-tert-butyl-1-phenyl-1H-pyrazol-5-yl) imino]methyl] phenol: Experimental and DFT computational studies. *Journal of Molecular Structure*, 1150: 366-373.

Samir Bondock, Ramy Rabie, Hassan A. Etman, Ahmed A. Fadda (2008). Synthesis and antimicrobial activity of some new heterocycles incorporating antipyrine moiety. *European Journal of Medicinal Chemistry*, 43:2122-2129.

Samir Bondock, Walid Fadaly, Mohamed A. Metwally. (2010). Synthesis and antimicrobial activity of some new thiazole, thiophene and pyrazole derivatives containing benzothiazole moiety. *European Journal of Medicinal Chemistry*, 45: 3692-3701.

Saugata Konar, Atanu Jana, Kinsuk Das, Sangita Ray, Sudipta Chatterjee, James A. Golen, Sayed, A. Z., Aboul-Fetouh, M. S. & Nassar, H. S. (2011). Synthesis, biological activity and dyeing performance of some novel azo disperse dyes incorporating pyrazolo [1, 5-a] pyrimidines for dyeing of polyester fabrics. *Journal of Molecular Structure*, 1010: 146-151.

Schiff, H. (1864). Mittheilungen aus dem Universitätslaboratorium in Pisa: eine neue Reihe organischer Basen. *Justus Liebigs Annalen der Chemie*, 131(1): 118-119.

Schmatz, D. M., Romancheck, M. A., Pittarelli, L. A., Schwartz, R. E., Fromling, R. A., Nollstadt, K. H. & Turner, M. J. (1990). Treatment of *Pneumocystis carinii* pneumonia with 1, 3-beta-glucan synthesis inhibitors. *Proceedings of the National Academy of Sciences*, 87(15): 5950-5954.

Sevil Ozkinali, Mahmut Gur, Nesrin Sener, Sati Alkin, M. Serdar Çavus (2018). Synthesis of new azo schiff bases of pyrazole derivatives and their spectroscopic and theoretical investigations. *Journal of Molecular Structure*, 1174: 74-83.

Shah, N. S., A. Biradar, S.I. Habib, J. Dhole, M. Baseer, P. Kulkarni (2011). Synthesis and antimicrobial studies of some novel pyrazolines. *Der Pharmaceutics Chemistry*, 3:167- 171.

Shehab A. Sallam (2004). Synthesis, characterization and thermal decomposition of copper (II), nickel (II) and cobalt (II) complexes of 3-amino-5-methylpyrazole Schiff-bases. *Transition Metal Chemistry*, 30:341-351.

Shelke, S. N., Mhaske, G. R., Bonifacio, V. D. & Gawande, M. B. (2012). Green synthesis and anti-infective activities of fluorinated pyrazoline derivatives. *Bioorganic & Medicinal Chemistry Letters*, 22(17): 5727-5730.

Shridhar Malladi, Arun M. Isloor, Shrikrishna Isloor, D.S. Akhila, Hoong-Kun Fun (2013). Synthesis, characterization and antibacterial activity of some new pyrazole based schiff bases. *Arabian Journal of Chemistry*, 6: 335-3340.

Song, M. X., Zheng, C. J., Deng, X. Q., Sun, L. P., Wu, Y., Hong, L. & Piao, H. R. (2013). Synthesis and antibacterial evaluation of rhodanine-based 5-aryloxy pyrazoles against selected methicillin resistant and quinolone-resistant *Staphylococcus aureus* (MRSA and QRSA). *European Journal of Medicinal Chemistry*, 60: 376-385.

Sumangala V., Boja Poojary, Chidananda N., Arulmoli T. and Shalini Shenoy (2012). Synthesis and biological evaluation of 2,4-disubstituted-[1,3]-thiazole. *Journal of Chemical and Pharmaceutical Research*, 4(12):4979-4987.

Susmita Mandal, Mousumi Das, Piu Das, Amalesh Samanta, Ray J. Butcher, Manan Saha, Ibrahim A. Alswaidan, Lydia Rhyman, Ponnadurai Ramasami, Nitis Chandra Saha (2018). Synthesis, characterization, DFT and antimicrobial studies of transition metal ion complexes of a new schiff base ligand, 5-methylpyrazole-3yl-N-(2-hydroxyphenyl amine) methylene imine, (MPzOAP). *Journal of Molecular Structure*, 1178:100-111.

Susmita Mandal, Ratan Sadhukhan, Utpal Ghosh, Soumik Mandal, Manan Saha, Ray J. Butcher & Nitis Chandra Saha (2016). Synthesis and characterization of two Cu (II) complexes with a new pyrazole-based Schiff base ligand: Crystallography, DNA interaction and antimicrobial activity of Ni (II) and Cu (II) complexes, *Journal of Coordination Chemistry*, 69(10): 1618-1634.

Taghrid Hafez, Souad A. Osman, Hisham Abdallah A. Yosef, Amira S. Abd El-ALL, Aashraf S. Hassan, Abdallah A. El-Sawy, Mohamed Abdallah, Mahmoud Youns. (2013). Synthesis, structural Elucidation, and In vitro Antitumor activities of some pyrazolopyrimidines Schiff bases derived from 5-amino-3-(aryl amino)-1H-pyrazole-4-carboxamides. *Science Pharmaceutical*, 81: 339-357.

Tarak Nath Mandal, Somnath Roy, Anil Kumar Barik, Samik Gupta, Ray J. Butcher, Susanta Kumar Kar (2008). Synthesis and structural characterization of copper (II) and vanadium(V)complexes of pyridyl/pyrimidyl-pyrazole derived Schiff base ligands - Metal specific adjustment of ligand binding mode. *Polyhedron*, 27: 3267-3274.

Thakar. A.S, K. S. Pandya, K. T. Joshi, A. M. Pancholi (2011). Synthesis, characterization and antibacterial activity of novel schiff bases derived from 4-Phenyl-2-amino thiazole and their Mn(II), Fe(II), Co(II), Ni(II) and Cu(II) metal complexes. *European Journal of Chemistry*, 8(4): 1556-1565.

Tripodi, A., Padovan, L., Veena, C., Scalabrinio, E., Testa, S. & Peyvandi, F. (2015). How the direct oral anticoagulant apixaban affects thrombin generation parameters. *Thrombosis Research*, 135(6): 1186-1190.

Uslaner, J. M., Parmentier-Batteur, S., Flick, R. B., Surles, N. O., Lam, J. S., McNaughton, C. H. & Hutson, P. H. (2009). Dose-dependent effect of CDPPB, the mGluR5 positive allosteric modulator, on recognition memory is associated with GluR1 and CREB phosphorylation in the prefrontal cortex and hippocampus. *Neuropharmacology*, 57(5-6): 531-538.

Vadiyala Naveen Reddy, Lingala Yamini, Yerrabelli Jayaprakash Rao, Chitneni Prasad Rao (2017). Synthesis of pyrazole-4-carbaldehyde derivatives for their antifungal activity. *Medicinal Chemistry Research*, 26:1664-1674.

Valeria La Pietra, Giuseppe La Regina, Antonio Coluccia, Valeria Famiglini, Sveva Pelliccia, Batya Plotkin, Hagit Eldar-Finkelman, Andrea Brancale, Carlo Ballatore, Alex Crowe, Kurt R Brunden, Luciana Marinelli, Ettore Novellino, and Romano Silvestri (2013). Design, synthesis, and biological evaluation of 1-Phenyl pyrazolo [3,4-e] pyrrolo [3,4-g] indolizine-4,6(1H,5H)-diones as New glycogen synthase Kinase-3# Inhibitors. *Journal of Medicinal Chemistry*, 56(24):10066-10078.

Vijesh, A. M., Isloor, A. M., Shetty, P., Sundershan, S. & Fun, H. K. (2013). New pyrazole derivatives containing 1, 2, 4-triazoles and benzoxazoles as potent antimicrobial and analgesic agents. *European Journal of Medicinal Chemistry*, 62: 410-415.

Vovk. M. V, A.V. Bol but, D.M. Volochnyuk, and A.M. Pinchuk. (2003). Synthesis of N-(5-pyrazolyl) schiff bases derived from Aryl trifluoromethyl ketones. *Russian Journal of Organic Chemistry*, 40: 63-66.

Wu, T. W., Zeng, L. H., Wu, J. & Fung, K. P. (2002). Myocardial protection of MCI-186 in rabbit ischemia-reperfusion. *Life Sciences*, 71(19): 2249-2255.

Xing, C., Li, X., Zhu, S., Zhao, J. & Zhu, S. (2006). Synthesis and hetero-Diels-Alder reactions of (E)- α -perfluoroalkanesulfonyl- α , β -unsaturated ketones. *Tetrahedron letters*, 47(28): 4951-4955.

Xu, L., Wang, J., Xiao, B., Yang, J., Liu, Y. & Huang, R. (2012). Preparation and characterization of a novel polymorph of indiplon, Form α . *Bioorganic & Medicinal Chemistry Letters*, 22(2): 963-968.

Yan, J., Wang, Q., Wei, T. & Fan, Z. (2014). Recent advances in design and fabrication of electrochemical supercapacitors with high energy densities. *Advanced Energy Materials*, 4(4): 1300816.

Zeitsch, K. J. (2000). *The Chemistry and Technology of Furfural and its Many By-Products* (Vol. 13). Elsevier.

Zeynizadeh, B. & Setamidideh, D. (2005). Water as a green solvent for fast and efficient reduction of carbonyl compounds with NaBH₄ under microwave irradiation. *Journal of the Chinese Chemical Society*, 52(6): 1179-1184.

