

ADVANCES IN NEUROMUSCULAR REHABILITATION STRATEGIES FOR STROKE SURVIVORS: A NARRATIVE REVIEW

Iram Hassan^{*1}, Umera Iftikhar², Faryal Baseerat Dar³, Aima Mushtaq⁴, Ahsan Hanif⁵
Samara Shaukat⁶

^{*1,4}Student Superior University, Lahore

²Assistant Professor Superior University, Lahore

³ Student Afro Asian Institute

^{5,6} Lecturer Superior University, Lahore

^{*1}hassaniramm808@gmail.com, ²umera.iftikhar@superior.edu.pk, ³faryalbaseerat@gmail.com,
⁴aimamushtaq52@gmail.com, ⁵ahsanhanif501@gmail.com, ⁶samara.shaukat@superior.edu.pk

DOI: <https://doi.org/10.5281/zenodo.17491003>

Keywords

Neuromuscular Rehabilitation,
Stroke Survivors, Motor Recovery,
Hemiparesis, Task-Specific
Training, Neuromuscular
Electrical Stimulation (NMES),
Neurorehabilitation

Article History

Received: 28 August 2025

Accepted: 19 October 2025

Published: 31 October 2025

Copyright @Author

Corresponding Author: *

Iram Hassan

Abstract

Objectives

This narrative review seeks to consolidate existing global research on neuromuscular rehabilitation strategies for stroke survivors, emphasizing therapeutic advancements, specific clinical populations, and the impact of these interventions on functional outcomes and healthcare service provision.

Data Sources

A search was conducted in electronic databases such as PubMed, Cochrane Library, PEDro, MEDLINE, and CINAHL for pertinent articles published between 2000 and 2023.

Review Methods

Research articles were selected based on their significance to neuromuscular rehabilitation following a stroke and their publication in peer-reviewed journals. A content analysis was conducted on randomized controlled trials (RCTs), systematic reviews, and observational studies published in English to identify essential findings concerning intervention strategies, target populations, and healthcare outcomes.

Results

A total of sixteen studies fulfilled the inclusion criteria. Commonly employed neuromuscular interventions in adult stroke populations included neuromuscular electrical stimulation (NMES), robotic-assisted therapy, task-specific training, and sensorimotor integration. The reported advantages encompassed enhanced motor control, improved functional mobility, and increased patient satisfaction. Nevertheless, the variability in methodologies restricted the generalizability of the findings.

Conclusions

Neuromuscular rehabilitation shows considerable potential in improving motor recovery after a stroke, especially in adults experiencing hemiparesis. However, it is essential to conduct future high-quality research to develop standardized, evidence-based protocols.

INTRODUCTION

Stroke ranks among the primary contributors to long-term disability globally, frequently leading to neuromuscular impairments such as hemiplegia, spasticity, and coordination difficulties (1). Neuromuscular rehabilitation seeks to promote motor recovery by employing methods that focus on enhancing neural plasticity and reconditioning muscles (2).

Physiotherapists play a key role in stroke management, employing evidence-based interventions that leverage the principles of neuroplasticity and motor learning (3). Recent innovations such as robotic therapy, NMES, and task-specific training have expanded the scope and impact of rehabilitation in stroke survivors (4).

Identification of Literature

A comprehensive literature review was performed utilizing the keywords: 'stroke,' 'neuromuscular rehabilitation,' 'robotic therapy,' 'task-specific training,' 'electrical stimulation,' and 'sensorimotor integration.' Research articles published from 2000 to 2023 were sourced from databases including PubMed, PEDro, Cochrane, CINAHL, and MEDLINE.

Inclusion criteria:

- Neuromuscular impairments associated with stroke.
- Utilization of physiotherapy-driven neuromuscular interventions.
- Full-text articles in English that have undergone peer review.

The articles were evaluated using the PEDro scale and CASP assessment tools.

Findings

Patient Populations

The majority of research focused on adults between the ages of 45 and 80 who have experienced chronic or subacute ischemic or hemorrhagic strokes (5). The presence of hemiparesis and compromised balance were common inclusion criteria (6).

Clinical Conditions Managed

Neuromuscular rehabilitation addresses deficiencies including:

- Muscle weakness
- Abnormal muscle tone (spasticity)
- Impaired coordination
- Reduced postural control and gait dysfunction (7).

Interventions Used

1. Neuromuscular Electrical Stimulation (NMES)

Neuromuscular Electrical Stimulation uses electric currents to make paralyzed or weak muscles contract. This treatment helps improve muscle strength, increase the ability to control movement on your own, and stop muscles from shrinking in the affected limb after a stroke (8). NMES is often used for shoulder subluxation, where electric stimulation of the supraspinatus and posterior deltoid muscles helps decrease the severity of the subluxation and the pain that comes with it (9). It is also helpful for foot drop, helping to lift the foot properly while walking and making walking faster and safer. How it works is that NMES can make the brain more active, which helps the brain change and improve movement abilities. Usually, each treatment session lasts between 20 to 60 minutes, and it is done 3 to 5 times a week, depending on how the patient feels and what their goals are (10).

2. Task-Specific Training (TST)

Task-specific training is a type of therapy that involves repeating useful activities to help someone become more independent in daily tasks. It is based on the idea that practicing specific skills over and over helps the brain adapt and improve. The method focuses on doing activities that have a clear purpose, repeating them regularly, and doing them in real-life situations to encourage the brain to reorganize itself (11). TST has shown big improvements in how well someone can use their arm, reach for things, and move around, especially when used along with constraint-induced movement therapy for the affected arm. For walking recovery, TST includes doing walking exercises, climbing stairs, or moving around obstacles, which helps improve movement and balance. Studies show that TST works best when done a lot and regularly

over several weeks, with each person's plan changing based on how they are doing (12).

3. Robotic-Assisted Therapy

Robotic-assisted therapy uses machines to help move weak limbs, offering lots of repeated practice focused on specific tasks. Tools like Lokomat are used for walking recovery and Armeo for arm training, allowing patients to do many movements with accurate control and instant feedback (13). Clinical studies show that using robots in therapy helps people walk better, control their arms more effectively, and have more stamina, especially for those who had a stroke not too long ago. These robotic systems also let doctors measure how people move, which helps them track improvements and change the therapy as needed. When robot therapy is used along with regular treatments, it can lead to better results by supporting both movements that people do on their own and those that are helped by the robot (14).

4. Sensorimotor Integration Therapy

Sensorimotor Integration Therapy helps improve how the body senses and responds to movement. It works by strengthening the senses that help you know where your body is, how it feels, and your balance. For people who have had a stroke, problems with these senses can make it harder to move smoothly, stay balanced, and coordinate movements (15). Sensorimotor training usually involves activities that require carrying weight, balancing on uneven surfaces, and repeating movements that help the body understand where its parts are. This type of training helps the brain better connect what the senses feel with how the body moves. Studies show that these exercises can improve balance, lower the chance of falling, and make everyday tasks easier. Combining sensorimotor training with other types of training, like task-focused or robotic exercises, can help the brain adapt and lead to better overall results (16).

Discussion

Neuromuscular rehabilitation strategies provide various pathways to enhance functional recovery following a stroke. Research consistently shows that early, intensive, and task-oriented interventions result in improved motor outcomes (17).

Electrical Stimulation:

NMES has demonstrated significant advantages in alleviating spasticity and enhancing muscle strength when combined with functional training (18).

Robotic Therapy:

Robotic systems provide secure and scalable rehabilitation solutions. Although they demonstrate potential, their elevated costs and restricted accessibility hinder their widespread adoption (19).

Task-Specific Training:

Among the most well-supported methods, TST improves motor control by promoting neuroplastic alterations in injured regions of the brain (20). Although the results are favorable, variations in study design, sample sizes, and outcome measures restrict the capacity to apply findings universally across different populations.

Limitations of Research

Limitations of the reviewed studies include:

- Limited sample sizes, diverse intervention protocols.
- Brief follow-up periods.
- Restricted application of objective neurophysiological outcome measures.
- Language restrictions and exclusion of grey literature may have omitted relevant studies.

Future Research Directions

To strengthen the field, future studies should:

- Implement extensive randomized controlled trials (RCTs) adhering to standardized protocols.
- Include long-term follow-up to evaluate the sustainability of motor improvements.
- Employ neuroimaging techniques to investigate the associated neural alterations.
- Assess the cost-effectiveness of conventional rehabilitation methods in comparison to technology-assisted approaches.
- Investigate hybrid rehabilitation models that integrate home-based practices with tele-rehabilitation technologies.

Conclusion

Neuromuscular rehabilitation has shown significant advantages in the recovery process following a stroke,

especially for adult survivors experiencing motor deficits. Techniques such as neuromuscular electrical stimulation (NMES), robotic-assisted therapy, and task-oriented training provide quantifiable enhancements in mobility, autonomy, and overall quality of life. Nonetheless, additional rigorous research is required to confirm and standardize these methods for wider application in clinical settings.

REFERENCES

1. Feigin VL, Vos T, Alahdab F, Amit AML, Bärnighausen TW, Beghi E, et al. Burden of neurological disorders across the US from 1990-2017: a global burden of disease study. *JAMA neurology*. 2021;78(2):165-76.
2. Dobkin BH. Rehabilitation after stroke. *New England Journal of Medicine*. 2005;352(16):1677-84.
3. Arena R, Bates B, Cherney LR, Cramer SC, Deruyter F, Eng JJ, et al. *AHA/ASA Guideline*. 2016.
4. Pollock A, Farmer SE, Brady MC, Langhorne P, Mead GE, Mehrholz J, et al. Cochrane overview: interventions for improving upper limb function after stroke. *Stroke*. 2015;46(3):e57-e8.
5. Hill VA, Towfighi A, editors. *Modifiable risk factors for stroke and strategies for stroke prevention*. Seminars in neurology; 2017: Thieme Medical Publishers.
6. Bernhardt J, Hayward KS, Dancause N, Lannin NA, Ward NS, Nudo RJ, et al. A stroke recovery trial development framework: consensus-based core recommendations from the second stroke recovery and rehabilitation roundtable. *International Journal of Stroke*. 2019;14(8):792-802.
7. Klijn CJ, Paciaroni M, Berge E, Korompoki E, Kőrv J, Lal A, et al. Antithrombotic treatment for secondary prevention of stroke and other thromboembolic events in patients with stroke or transient ischemic attack and non-valvular atrial fibrillation: a European Stroke Organisation guideline. *European Stroke Journal*. 2019;4(3):198-223.
8. Haque F. Effectiveness of Functional Electrical Stimulation (FES) on upper limb motor functional recovery in people who experienced a stroke: Bangladesh Health Professions Institute, Faculty of Medicine, the University ...; 2023.
9. Lee J-H, Baker LL, Johnson RE, Tilson JK. Effectiveness of neuromuscular electrical stimulation for management of shoulder subluxation post-stroke: a systematic review with meta-analysis. *Clinical rehabilitation*. 2017;31(11):1431-44.
10. Everaert DG, Thompson AK, Chong SL, Stein RB. Does functional electrical stimulation for foot drop strengthen corticospinal connections? *Neurorehabilitation and neural repair*. 2010;24(2):168-77.
11. Langhorne P, Coupar F, Pollock A. Motor recovery after stroke: a systematic review. *The Lancet Neurology*. 2009;8(8):741-54.
12. Winstein CJ, Wolf SL, Schweighofer N. Task-oriented training to promote upper extremity recovery. Stein J, Harvey RL, Macko RF, Winstein CJ, Zorowitz Rd *Stroke Recovery and Rehabilitation* New York: Demos Medical. 2008:267-90.
13. Lo AC, Guarino PD, Richards LG, Haselkorn JK, Wittenberg GF, Federman DG, et al. Robot-assisted therapy for long-term upper-limb impairment after stroke. *New England Journal of Medicine*. 2010;362(19):1772-83.
14. Mehrholz J, Pohl M, Platz T, Kugler J, Elsner B. Electromechanical and robot-assisted arm training for improving activities of daily living, arm function, and arm muscle strength after stroke. *Cochrane database of systematic reviews*. 2018(9).
15. Carey LM. Review on somatosensory loss after stroke. *Critical Reviews™ in Physical and Rehabilitation Medicine*. 2017;29(14).
16. De Bruyn N, Essers B, Thijs L, Van Gils A, Tedesco Triccas L, Meyer S, et al. Does sensorimotor upper limb therapy post stroke alter behavior and brain connectivity differently compared to motor therapy? Protocol of a phase II randomized controlled trial. *Trials*. 2018;19(1):242.
17. Veerbeek JM, van Wegen E, van Peppen R, Van der Wees PJ, Hendriks E, Rietberg M, et al. What is the evidence for physical therapy poststroke? A

- systematic review and meta-analysis. PloS one. 2014;9(2):e87987.
18. Sabut SK, Sikdar C, Mondal R, Kumar R, Mahadevappa M. Restoration of gait and motor recovery by functional electrical stimulation therapy in persons with stroke. Disability and rehabilitation. 2010;32(19):1594-603.
 19. Louie DR, Eng JJ. Powered robotic exoskeletons in post-stroke rehabilitation of gait: a scoping review. Journal of neuroengineering and rehabilitation. 2016;13:1-10.
 20. French MA, Morton SM, Reisman DS. Use of explicit processes during a visually guided locomotor learning task predicts 24-h retention after stroke. Journal of Neurophysiology. 2021;125(1):211-22.

