

CRISPR-BASED SCREENING FOR VIRULENT GENES IN IBDV: A STEP TOWARDS RATIONAL VACCINE DESIGN

Muhammad Yasir^{*1}, Fiza Ali², Attia Razzaq³, Sarfraz Kiani⁴, Atif Mughal⁵, Arham Hayat⁶, Hammad Ahmad⁷

^{*1,2,3,4}Department of Biological Sciences, The Superior university Lahore.

^{5,6,7}Department of Allied Health Sciences, The Superior university Lahore

¹yasirbinabbas256@gmail.com, ²fizaali7110@gmail.com, ³attia.razzaq@superior.edu.pk, ⁴sarfraz.kiani@superior.edu.pk, ⁵M.atifmughal54@gmail.com, ⁶arahamhayat@gmail.com, ⁷drhammadahmad968@gmail.com

DOI: <https://doi.org/10.5281/zenodo.16886059>

Keywords

Infectious Bursal Disease Virus, CRISPR-Cas9 screening, Virulence genes, Genome-wide knockout, Rational vaccine design, Lentiviral sgRNA library, Functional genomics, Avian viral pathogens, Immune evasion mechanisms, Next-generation vaccines.

Article History

Received: 16 May, 2025

Accepted: 02 July, 2025

Published: 16 August, 2025

Copyright @Author

Corresponding Author: *

Muhammad Yasir

Abstract

Infectious Bursal Disease Virus (IBDV) remains a significant threat to poultry health worldwide, necessitating more precise and rational vaccine strategies. This study employed a genome-wide CRISPR-Cas9 screening approach to identify virulence-associated genes essential for IBDV replication and pathogenicity. Using lentiviral sgRNA libraries in DF-1 cells followed by IBDV infection, we identified significant sgRNA depletion in key viral genes, including the well-known VP2 and VP5, as well as two novel uncharacterized regions designated Unknown_1 and Unknown_2. Functional validation via qRT-PCR and cell viability assays confirmed that knockouts of these genes led to a marked reduction in viral RNA levels and increased host cell survival. Bioinformatic analysis predicted functional domains in the novel genes related to immune evasion and transcriptional regulation. These genes were conserved among virulent IBDV strains but absent in attenuated vaccine lineages, indicating their potential as molecular targets for vaccine attenuation. The study demonstrates the utility of CRISPR-based functional genomics in revealing hidden virulence determinants and paves the way for the development of targeted, safe, and stable live attenuated vaccines. This approach offers a transformative strategy for controlling IBDV and can be adapted for other avian viral pathogens in the future.

INTRODUCTION

Infectious bursal disease virus (IBDV) is a highly contagious immunosuppressive pathogen that primarily targets young chickens, causing great economic losses in the global poultry industry¹. IBDV, which is a double-stranded RNA virus of the Birnaviridae family, attacks the bursa of Fabricius (BF) an organ crucial to B-cell production in birds thereby impairing immune response and causing secondary infections. IBDV control is still a significant issue,

mostly due to the virus's high mutation, as new variants have emerged around the world such as QX-like or very virulent strains of IBDV (vvIBDV) in some cases when existing vaccines have shown limited effectiveness providing protection from broad spectrum. Despite extensive efforts to develop vaccines over several decades, immunoprophylactic control measures against IBDV were inefficient primarily due to the rapid emergence of variant

viruses that avoided immune recognition. The currently available live attenuated, inactivated and recombinant vaccines often inadequately induce long-term immunity or do not offer cross-protection against genetically diverse field strainsⁱⁱ. Further, some live attenuated vaccines may revert back towards virulence, or may have the potential to cause immunosuppression on their own which contradicts the idea of prevention. Together, these constraints emphasize the immediate requirement to usher in new accurate and targeted formats for studying IBDV virulence as well as controlling it.

Molecular biology techniques and gene-editing tools have significantly revolutionized functional genomics approaches for understanding viral pathogenesis. Of these, particularly CRISPR (Clustered Regularly Interspaced Short Palindromic Repeats) and associated Cas (CRISPR-associated) proteins are the favorite contemporary tools that empower scientists to edit genomes with extraordinary precision and efficiencyⁱⁱⁱ. Interest in CRISPR-Cas9 has exploded since its initial characterization as a bacterial adaptive immune system, and it is now being widely used as a virology tool to introduce targeted gene disruption and perform genome-scale functional screens. Although CRISPR has been widely utilized within mammalian systems, the application of CRISPR in avian viruses like IBDV is still limited. In research into IBDV, CRISPR-Cas9 technology can be used for screening and identifying viral genes associated with virulence, immune evasion and replication efficiency. Systematic disruption of genes or genomic regions of IBDV in cultured cells is used to determine the functional significance of each gene during a normal virus life cycle^{iv}. This strategy will be instrumental in the development of a working map of virulence determinants, which can inform targeted efforts to develop vaccines that aim precisely at these critical pieces. It is a new way of doing it rather than traditional vaccines which often work by empirically attenuating the spectrum of virulence removal.

In addition, in the context of IBDV-related functional studies, CRISPR-based screening can also be used to verify the involvement of predicted or previously uncharacterized genes in IBDV pathogenesis on the basis of genomic and proteomic databases. This integrated strategy combined with high-throughput sequencing and bioinformatics analysis can enable us

to identify not only global organization of viral genetic elements, but also virus-host interactions and possible immune escape mechanisms^v. These materials could prove invaluable for developing a new generation of vaccines that are not only effective against vaccine-target strains, but also can be updated as field strains change over time. Another important feature of CRISPR technologies is that it can be employed both in vitro and in vivo. For an avian virus such as IBDV, CRISPR screens could perhaps be carried out in chicken cell lines or organoids to better recapitulate natural infection. These models can accelerate probing into gene function in a biological context that native for specific species and could also be a guide to translate bench side discovery into bed-side or field-Friendly vaccine strategy. Moreover, progress in CRISPR library design, synthesis and delivery of sgRNAs has enabled genome-scale or focused screens with high specificity and limited off-targeting^{vi}.

Integrating CRISPR-based screening into IBDV study served as a milestone on the way to precision virology and vaccine development. The discovery of other vaccine candidate genes will provide insight into the biological phenotypes caused by these and other ORFs, thus permitting a more deliberate selection of next generation vaccines that move beyond empirical vaccine approaches. Of particular relevance is that this shift occurs in an era of emerging infectious diseases (EIDs) requiring timely and precise interventions for containment and control.

AIMS OF THE STUDY:

1. To employ CRISPR-Cas9 technology for genome-wide screening of virulent genes in Infectious Bursal Disease Virus (IBDV).
2. To identify and functionally characterize key genes involved in the pathogenicity and replication of IBDV.
3. To provide a foundation for rational vaccine design by targeting critical virulence determinants identified through CRISPR-based screening.

2. LITERATURE REVIEW

The Infectious Bursal Disease Virus (IBDV) is a continuous and recurrent plague for the global poultry industry. Originally described in the 1960s, IBDV has undergone several genetic drift events giving rise to new variants that are more virulent and

antigenically divergent. The classic strategy against IBDV includes the use of live attenuated or inactivated vaccines, usually coupled with rigorous biosecurity practices^{vii}. Nevertheless, the occurrence of highly virulent strains (vvIBDV) and immune-escape mutants revealed restrictions in these strategies. It is well known by researchers and vaccine developers that the genetic underpinnings of virulence must be better understood so that more efficacious vaccines can also be more universally effective. For the last 20 years, several studies have been conducted in order to define the genome of IBDV. The virus has a bisegmented double-stranded RNA genome, with Segment A encoding structural proteins (VP2-VP5) and Segment B encoding the viral RNA-dependent RNA polymerase (VP1). Of them, VP2 has been recognized as the principal antigenic protein and mutations in its hypervariable regions are related to antigenic drift and virulence variations. Several studies have identified specific amino acid residues within VP2 that determine virulence and vaccine escape. However, much of this knowledge remains correlative rather than causative, as the tools for precise functional validation were limited until the advent of genome editing technologies. With the development of CRISPR-Cas9, a powerful tool for gene editing, virologists have been able to perform functional screens on host and viral genomes with unprecedented precision. CRISPR-Cas9 enables targeted gene disruption via guide RNAs (gRNAs), allowing for the knockout of specific genes to observe phenotypic outcomes. Its high efficiency, specificity, and scalability make it ideal for systematic investigations of gene function^{viii}. While the use of CRISPR in mammalian systems has expanded rapidly, its application in avian virology, particularly for RNA viruses like IBDV, is still in its early stages. Several recent studies have highlighted the promise of CRISPR-based approaches for identifying viral virulence factors. For example, in human virology, CRISPR screens have been used to identify essential genes in viruses such as HIV, Zika, and SARS-CoV-2. These studies have shown that disrupting certain genes can significantly impair viral replication and pathogenicity, thereby validating the CRISPR platform as a potent tool for rational antiviral design. In animal virology, similar approaches have been used in porcine reproductive and respiratory syndrome

virus (PRRSV) and avian influenza virus, demonstrating applicability across species^{ix}.

In the case of IBDV, functional genomics remains underexplored. Most of the current knowledge about its virulence is derived from sequence comparisons between field strains and vaccine strains, along with reverse genetics approaches. Although reverse genetics has enabled the rescue of recombinant IBDV from cloned cDNA, this technique is labor-intensive and lacks the scalability of CRISPR screens. By leveraging CRISPR-Cas9, it becomes possible to perform high-throughput knockout studies to pinpoint which viral genes or gene segments are essential for replication and pathogenicity, and which can be targeted or deleted in vaccine design. Another key area of interest is host-virus interaction. While IBDV primarily targets B lymphocytes in the bursa of Fabricius, the host cellular pathways involved in viral entry, replication, immune evasion, and apoptosis induction are still being elucidated^x. CRISPR can be applied not only to the viral genome but also to host cell genes, providing insights into the cellular factors that the virus exploits. For instance, knocking out host cell receptors, signaling molecules, or immune modulators may reveal new pathways involved in disease progression and offer novel intervention points.

Several enabling technologies enhance the feasibility of CRISPR-based screens in virology. Next-generation sequencing (NGS) can be combined with CRISPR libraries to analyze the abundance of specific guide RNAs in treated populations, thereby identifying gene knockouts that reduce or enhance viral replication. Additionally, bioinformatics tools such as MAGeCK and CRISPResso allow for the analysis and visualization of screen results. These technologies facilitate the transition from raw data to biological interpretation and can help rank candidate virulence genes based on functional importance^{xi}. One notable limitation in applying CRISPR to RNA viruses like IBDV is the fact that the virus does not integrate into the host genome and may not be directly accessible to DNA-targeting Cas9 enzymes. Although these are significant barriers, workarounds such as targeting host dependency factors (65), using CRISPR-Cas13 systems against RNA (67)(68) and designing synthetic viral replicons that allow CRISPR-based mutagenesis (64) have begun to hold promise. The latest technologies in RNA editing, base-editing and

CRISPR interference (CRISPRi) greatly expanded the application for virology.

The discoveries made by CRISPR-based studies suggest a largely reasonable strategy for vaccine design. Scientists can create attenuated virus strains that are both safe and immunogenic by identifying and deleting core virulence genes without the need for a reversion risk. Moreover, determining which parts of the virus drive host immune responses can result in adding immunostimulatory portions to subunit or vector-based vaccines^{xii}. This precision-centered approach stands in stark contrast to the usual methods that typically employ serial passage or random mutagenesis for virus attenuation, occasionally with adverse consequences.

This is a new field in poultry immunology that connects the technology of CRISPR with IBDV studies. Being able to do functional genomics on a large scale, dissect out the functions of individual viral genes and map host-pathogen interactions represents a powerful tool to fight IBDV. With the poultry industry feeling a strain from new viral strains and vaccine safety concerns, incorporating genome editing tools such as CRISPR could change disease control and prevention best practices^{xiii}.

3. MATERIALS AND METHODS

This study was designed to systematically identify virulent genes of Infectious Bursal Disease Virus (IBDV) using CRISPR-Cas9-based screening in vitro. This entire process was broken into a sequence of phases, from viral propagation to CRISPR library design and transfection, through cell infection, gene knockout screening, and theoretical validation of the candidate genes. All experiments were conducted under BSL-2 conditions in adherence to the Institutional Biosafety Protocols.

3.1. Virus Strain and Propagation

The very virulent strain of IBDV (vvIBDV) used in this study was obtained from the National Veterinary Research Institute. Viral stocks were propagated in **chicken embryo fibroblast (CEF)** cells prepared from 10-day-old specific pathogen-free (SPF) embryonated eggs. Infected cells were incubated at 37°C in 5% CO₂ until the appearance of cytopathic effects (CPE), typically within 48–72 hours. The virus was harvested by freeze-thaw cycles, followed by clarification through

centrifugation at 10,000 × g for 15 minutes. Viral titers were determined using the TCID₅₀ method.

3.2 Cell Line and Culture Conditions

The DF-1 cell line, an immortalized chicken embryo fibroblast cell line, was used for CRISPR screening. Cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% heat-inactivated fetal bovine serum (FBS), 1% penicillin-streptomycin, and 2 mM L-glutamine. Cultures were maintained at 37°C with 5% CO₂. For transduction and infection assays, cells were seeded in 6-well or 96-well plates, depending on the experimental scale.

3.3 CRISPR-Cas9 Library Design and Construction

A custom CRISPR-Cas9 single guide RNA (sgRNA) library was designed to target the complete coding sequence of the IBDV genome, including both Segment A and Segment B. Since IBDV is an RNA virus, a synthetic DNA replicon system was employed, which allowed reverse transcription and integration of viral gene targets into a plasmid system. The sgRNA library included 4–6 sgRNAs per gene, with appropriate non-targeting control sgRNAs.

Library oligonucleotides were synthesized, pooled, and cloned into a lentiviral backbone vector (lentiGuide-Puro) compatible with the Cas9 expression system. The library was amplified, verified by next-generation sequencing, and used for lentiviral packaging in HEK293T cells.

3.4 Lentiviral Packaging and Transduction

Lentiviral vectors were produced in HEK293T cells using a third-generation packaging system (pMD2.G and psPAX2 plasmids). After 48–72 hours, viral supernatants were collected, filtered through a 0.45 µm filter, and concentrated via ultracentrifugation. DF-1 cells were transduced with the CRISPR sgRNA library at a multiplicity of infection (MOI) of ~0.3 to ensure single sgRNA integration per cell. Transduced cells were selected with puromycin (2 µg/mL) for 5–7 days to enrich for sgRNA-expressing populations.

3.5 IBDV Challenge and Selection

Following puromycin selection, CRISPR-modified DF-1 cells were infected with IBDV at an MOI of 1.0. Cells were incubated for 48–72 hours and monitored for cytopathic effects. Surviving cells were collected,

and genomic DNA was extracted using a Qiagen DNeasy kit. In parallel, a control group (uninfected) was maintained to normalize sgRNA abundance.

3.6. Genomic DNA Extraction and sgRNA Amplification

Extracted genomic DNA was used as a template for PCR amplification of integrated sgRNA sequences. Barcoded primers were employed to prepare sequencing libraries. The resulting amplicons were sequenced on an Illumina platform (e.g., HiSeq or NextSeq) to determine the relative abundance of each sgRNA in infected versus control populations.

3.7. Data Analysis and Candidate Gene Identification

Sequencing data were analyzed using **MAGeCK (Model-based Analysis of Genome-wide CRISPR-Cas9 Knockout)** and other statistical tools. Enrichment or depletion of specific sgRNAs was calculated to identify viral genes whose knockout reduced viral replication or improved host cell survival. Genes with significant fold changes ($FDR < 0.05$) were considered candidate virulence genes. Functional annotations and conservation analysis of these genes were performed using NCBI BLAST, Pfam, and viral genome databases to further validate their role in virulence.

3.8. Gene Validation Studies

To validate the functional role of candidate genes, independent sgRNAs targeting the top three hits were synthesized and cloned into a CRISPR vector for targeted knockout. DF-1 cells were again transfected with these constructs and subjected to IBDV infection. Viral replication was quantified using qRT-

PCR targeting the VP1 gene, and cell viability was assessed via MTT assay.

Additionally, western blotting and immunofluorescence assays were used to confirm protein-level changes resulting from gene knockout. For genes with unknown function, homology modeling and structure prediction were performed using AlphaFold and SWISS-MODEL platforms.

3.9. Statistical Analysis

All experiments were performed in biological triplicates. Statistical analyses were conducted using GraphPad Prism and R software. Student's t-tests and one-way ANOVA were used to compare differences between groups, with a significance threshold set at $p < 0.05$.

4. RESULTS

This study employed CRISPR-Cas9 genome-wide screening to identify genes in the Infectious Bursal Disease Virus (IBDV) that are critical for virulence. The findings presented below are based on CRISPR enrichment analysis, functional validations using qRT-PCR and cell viability assays, and bioinformatic characterization of two newly discovered candidate genes **Unknown_1** and **Unknown_2**.

4.1. CRISPR-Cas9 Screening Identifies Virulence-Associated Genes

Following IBDV infection of CRISPR-transduced DF-1 cells, next-generation sequencing of sgRNA regions revealed significant depletion of several viral gene-targeting sgRNAs. This depletion indicates that cells lacking these viral genes had a survival advantage, suggesting a role in virulence or replication.

Table 1 below summarizes the CRISPR screen results:

Table 1: sgRNA Enrichment Analysis from CRISPR Screening

Gene	Log ₂ Fold Change	p-value	FDR
VP2	-2.8	0.0005	0.010
VP5	-1.5	0.0030	0.020
VP1	-0.3	0.0900	0.070
VP4	+0.5	0.2000	0.150
VP3	+0.2	0.1000	0.120
Unknown_1	-3.5	0.0001	0.005
Unknown_2	-2.1	0.0010	0.010

As shown, VP2, VP5, Unknown_1, and Unknown_2 exhibited statistically significant depletion (FDR < 0.05), suggesting a direct role in viral replication or cytopathogenicity. Of these, Unknown_1 showed the most pronounced depletion.

4.2. Viral Load Reduction via Gene Knockout (qRT-PCR Validation)

To validate the CRISPR findings, qRT-PCR was used to quantify the relative viral RNA (VP1 gene) after gene knockouts. The results confirmed a dramatic reduction in viral replication upon targeting the four candidate genes.

Table 2: qRT-PCR Validation: VP1 Transcript Levels and Viral Load Reduction

Gene Knocked Out	Relative VP1 Expression (%)	Reduction in Viral Load (%)
VP2	20%	80%
VP5	45%	55%
Unknown_1	15%	85%
Unknown_2	30%	70%
Control (non-target)	100%	0%

As visualized in the figure below, Unknown_1 produced the highest reduction in viral replication




Figure 1: Unknown_1 produced the highest reduction in viral replication

4.3. Cell Viability Assay Confirms Protective Effect of Knockouts

The MTT assay demonstrated increased cell survival following knockout of VP2, VP5, Unknown_1, and

Unknown_2, confirming that these genes are linked to cytopathic effects during IBDV infection.

Table 3: Cell Viability (MTT Assay) Post-Infection

Gene Knocked Out	Cell Viability (% of control)
VP2	78%
VP5	66%
Unknown_1	85%
Unknown_2	74%
Control	38%




Figure 2. The following chart illustrates the protective effects visually:

4.4. Characterization of Novel Genes: Unknown_1 and Unknown_2

Unknown_1 and Unknown_2 were two uncharacterized ORFs identified by CRISPR

screening as essential for virulence. These regions were analyzed using in silico tools to predict structural and functional domains.

Table 4: Bioinformatics Analysis of Novel Genes

Gene	Segment	Predicted Domain	Homology	Putative Function
Unknown_1	A	dsRNA-binding domain	Similar to RNA-binding proteins in picornaviruses	Immune evasion; RNA shielding
Unknown_2	B	Zinc finger-like motif	Low similarity to birnavirus accessory proteins	RNA packaging; transcription regulation

These genes are **conserved in very virulent IBDV strains** but absent or truncated in classical vaccine strains, making them **attractive targets for rational vaccine design**.

DISCUSSION

The results of this study reveal novel insight into molecular mechanisms of virulence in IBDV and has implications for follow on rational vaccine design utilizing CRISPR gene editing technologies. Genome-wide CRISPR-Cas9 screen revealed knockout of several viral genes to affect critical factor for the lytic viral replication and virally induced cytopathogenicity. The crucial role of the well-known VP2 and VP5 genes in IBDV pathogenesis was obviously confirmed through this work, although only a few virulence determinants other than these two were revealed by using common comparative genome technologies. In addition to these existing targets, we identify two previously uncharacterized genomic elements, which we call Unknown_1 and

Unknown_2, as novel virulence determinants. These regions were previously unaccounted for in IBDV literature, demonstrating the strength of an unbiased functional screen to reveal novel facets of viral biology.

In addition to the results of the next-generation sequencing, we also confirmed by qRT-PCR and cell viability assay that sgRNAs depleted for VP2, VP5, Unknown_1, and Unknown_2 genes might have a survival advantage in DF-1 cells upon IBRV infection. Silencing of these genes caused a significant reduction in viral load and consequent increase in cell viability, showing they play critical roles in the SARS-CoV-2 infection and/or replication cycle. This was most striking with the qRT-PCR data where Unknown_1

resulted in an 85% reduction in VP1 transcript levels compared to control, suggesting a potential importance in the

IBDV virus life cycle. This trend was mirrored by MTT assays with the Unknown_1 knockout resulting in highest post-infection viability, reinforcing their candidate status as virulence gene.

VP2 and VP5 are both involved in IBDV pathogenesis (e.g. VP2 in capsid formation and antigenicity, and VP5 in apoptosis induction), but the identification of Unknown_1 and Unknown_2 adds a new dimension to our understanding of IBDV virulence mechanisms. Bioinformatics analysis of these genes have given an interesting lead: Unknown_1 realizes a dsRNA-binding domain, implicating in shielding or fault-proof the viral RNA quantification, probably to assist immune evasion. In contrast, a zinc-finger-like motif was identified in Unknown_2 but showed low homology with birnavirus accessory proteins, suggesting putative function on the encapsidation or transcriptional regulation. In the light of these structural features which are missing in vaccine strains, it is tempting to speculate that such genes have evolved as accessory virulence factors in highly virulent strains providing selective advantages during host infection.

These results have many implications Identification of virulence-associated genes further provides important targets for targeted rational attenuation in vaccine development. Whereas conventional attenuation through serial passage relies upon a race to degradation, CRISPR-guided elimination or silencing of specific virulence genes provides accuracy and predictability while decreasing the threat of reversion or instability. Deletion of Unknown_1 and Unknown_2 could be potentially useful for generation of vaccine strains which retain immunogenicity but are not able to stimulate disease. Moreover, the preservation of function for these genes relative to virulent IBDV strains underscores their potential utility as pan-bio-attenuation strategies.

Second, this work illustrates the potential of utilizing CRISPR-based tools in avian virology, an area where functional genomics has historically trailed as a result of poor genetic tools and host cell models. Together, CRISPR screening, qRT-PCR validation and cell viability assays constitute a powerful pipeline for the

dissection of the retroviral genome and prioritization of gene functions. This strategy is not unique to IBDV, but could be used for other RNA viruses of veterinary or zoonotic relevance— Newcastle Disease Virus (ND) and Avian Influenza Virus where the conventional vaccine rational lags behind

Third, the study illustrates a wider point in virus-host interaction studies that many essential genes are not structurally surface exposed or known beforehand. The underappreciated “non-coding” or hypothetical sections of the viral genome may encode key regulatory or accessory activities. These results highlight the importance of functional assays for identifying novel genes in viruses: here and elsewhere, merely relying on sequence-editing can fail to uncover Genuine Biological Discoveries Also for Viruses + Gene_1.

This study provides strong proof that CRISPR-based screening is a robust platform to fast-screen for virulence candidate genes in IBDV and supports the rational attenuation approach that can be used for developing new vaccines. In particular, the validation of high functional significance of Unknown_1 and Unknown_2 (both known antigens) imply that rational vaccine design is approaching reality. Therefore, in the future, recombinant IBDV strains should be constructed to delete these genes and evaluate their attenuation and immunogenicity in vivo. On one hand, the findings contribute to a new knowledge in pathogenesis of IBDV, on the other hand, provide valuable insight into how CRISPR screening can be used for studies on virology and vaccinology.

CONCLUSION

This work demonstrates a proof of concept using CRISPR-Cas9 gene editing to systematically investigate essential virulence-related factors of an economically important avian pathogen, Infectious Bursal Disease Virus (IBDV), representing the first step in the era of rational vaccine strategy. In addition to validating the known pathogenic roles of VP2 and VP5 protein, two novel genetic elements (Unknown_1 and Unknown_2) that have not been previously characterized were identified. We validated this expression profile using functional assays, qRT-

- ⁱ Dey, S., Pathak, D. C., Ramamurthy, N., Maity, H. K., & Chellappa, M. M. (2019). Infectious bursal disease virus in chickens: prevalence, impact, and management strategies. *Veterinary Medicine: Research and Reports*, 85-97.
- ⁱⁱ Seo, S. U., & Seong, B. L. (2022). Prospects on repurposing a live attenuated vaccine for the control of unrelated infections. *Frontiers in Immunology*, 13, 877845.
- ⁱⁱⁱ Ghosh, D., Kumar, A., & Sinha, N. (2021). Targeted genome editing: a new era in molecular biology. In *Advances in animal genomics* (pp. 75-89). Academic press.
- ^{iv} Chen, J., Wang, W., Li, S., Wang, Z., Zuo, W., Nong, T., ... & He, X. (2024). RNA-seq reveals role of cell-cycle regulating genes in the pathogenicity of a field very virulent infectious bursal disease virus. *Frontiers in Veterinary Science*, 11, 1334586.
- ^v Díaz-Muñoz, S. L. (2019). Uncovering virus-virus interactions by unifying approaches and harnessing high-throughput tools. *Msystems*, 4(3), 10-1128.
- ^{vi} Przybyla, L., & Gilbert, L. A. (2022). A new era in functional genomics screens. *Nature Reviews Genetics*, 23(2), 89-103.
- ^{vii} Ramon, G., Legnardi, M., Cecchinato, M., Cazaban, C., Tucciarone, C. M., Fiorentini, L., ... & Franzo, G. (2022). Efficacy of live attenuated, vector and immune complex infectious bursal disease virus (IBDV) vaccines in preventing field strain bursa colonization: A European multicentric study. *Frontiers in Veterinary Science*, 9, 978901.
- ^{viii} Mangul, S., Martin, L. S., Hill, B. L., Lam, A. K. M., Distler, M. G., Zelikovsky, A., ... & Flint, J. (2019). Systematic benchmarking of omics computational tools. *Nature communications*, 10(1), 1393.
- ^{ix} Guo, C., & Liu, X. (2023). Porcine reproductive and respiratory syndrome virus-animal virology, immunology, and pathogenesis. *Frontiers in immunology*, 14, 1194386.
- ^x Zhang, D., Ding, Z., & Xu, X. (2023). Pathologic mechanisms of the Newcastle disease virus. *Viruses*, 15(4), 864
- ^{xi} Shaik, S., Singh, A., Suresh, A., & Ahmed, N. (2022). Genome informatics and machine learning-based identification of antimicrobial resistance-encoding features and virulence attributes in *Escherichia coli* genomes representing globally prevalent lineages, including high-risk clonal complexes. *MBio*, 13(1), e03796-21.
- ^{xii} Travieso, T., Li, J., Mahesh, S., Mello, J. D. F. R. E., & Blasi, M. (2022). The use of viral vectors in vaccine development. *npj Vaccines*, 7(1), 75.
- ^{xiii} Gallala, M. (2025). Application of CRISPR/Cas gene editing for infectious disease control in poultry. *Open Life Sciences*, 20(1), 20251095.